



88071825

QE
104
.P56
646
1983

PHASE I

Geology, Energy and Mineral (GEM) Resource
Evaluation of the Pioneer Mountains GRA,
including the Appendicitus Hill (31-14),
White Knob Mountains (31-17), Little Wood
River (53-4), and Friedman Creek (53-5)
Wilderness Study Areas

Bureau of Land Management
Contract No. YA-553-CT2-1039

By:

Greg Fernette
Joel Stratman

Contributors:

G. Webster
D. Blackwell
C.G. Bigelow
R.S. Fredericksen
J. Bressler
W.C. Jones
E.F. Evoy
J.F. McQuat

BLM Library
D-553A, Building 50
Denver Federal Center
P. O. Box 25047
Denver, CO 80225-0047

ANCHORAGE, ALASKA
OCTOBER 1983

WGM INC.
MINING AND GEOLOGICAL CONSULTANTS

50
D-452A, Ball
Library
Denver, CO 80202-0012


EXECUTIVE SUMMARY

The Pioneer Mountains GRA is located east of Ketchum in south-central Idaho. The area includes four BLM Wilderness Study Areas: Appendicitus Hill (31-14), White Knob Mountains (31-17), Little Wood River (53-4) and Friedman Creek (53-5). The GRA is in the Idaho Falls and Shoshone BLM districts.

There is good geologic data available on the west part of the Pioneer Mountains GRA which includes WSAs 53-4 and 53-5. The data on the east part of the GRA, including WSAs 31-14 and 31-17, is less detailed. The evaluation of oil and gas potential was greatly aided by Nance Petroleum Company who allowed examination of propriety geological and geophysical data from the east part of the area.


The bedrock in the Pioneer Mountains GRA consists of a thick sequence of sedimentary rocks which have been thrust faulted and folded into complex structures. The older rocks are overlain in much of the GRA by younger volcanic rocks. Locally, the sediments are intruded by small igneous stocks. The geophysical data for the area also indicates that several other subsurface intrusive bodies are present.

The Pioneer Mountains GRA includes all or portions of the Copper Basin, Little Wood River, and Lava Creek Mining districts. These areas were active in the late 1800s and early 1900s and produced silver, lead, zinc, copper, gold and barite. There has been intermittent mineral exploration in these district since active mining ceased. As of June 7, 1982 there were over 1,200 mining claims in the area. Much exploration activity is centered on



the Muldoon area just north of WSA 53-5. Because of this activity field work was done in WSA 53-5 as part of this evaluation. The work shows that favorable geologic structures and geochemical anomalies occur in the Friedman Creek WSA.

The Little Wood River and Friedman Creek WSAs are classified as moderately to highly favorable for metallic mineral resources. All of the WSA have moderate potential for oil and gas. The land classification for the WSAs is summarized in the accompanying table.



PIONEER MOUNTAINS GRA
SUMMARY OF GEM RESOURCES CLASSIFICATION

<u>Resource</u>	<u>Wilderness Study Areas</u>			
	<u>WSA 31-14</u>	<u>WSA 31-17</u>	<u>WSA 53-4</u>	<u>WSA 53-5</u>
1. Locatable				
a. Metallic Minerals	2A-3A	2A-3A	3C	4D
b. Uranium and Thorium	2A	2A	2B	2B
c. Non-metallic Minerals	2A-3A	2A-3A	3B	3C
2. Leasable				
a. Oil and Gas	3C	3C	3C	3C
b. Geothermal (L)	1B/2B	1B	1B	1B
Geothermal (H)	1B/2B	1B	1B	1B
c. Sodium-Potassium	2B	2B	2B	2B
d. Other (1)	2B	2B	2B	2B
3. Saleable	3B	3B	2B	2B

(1) Includes oil shale, asphalt, phosphate and bitumen.




TABLE OF CONTENTS

	<u>Page</u>
1.0 INTRODUCTION	1
1.1 Location	3
1.2 Population and Infrastructure	3
1.3 Basis of Report	3
1.4 Acknowledgements	6
2.0 GEOLOGY	7
2.1 Introduction	7
2.2 Physiography	7
2.3 Rock Units	9
2.4 Structural Geology and Tectonics	22
2.5 Paleontology	27
2.6 Historical Geology	30
3.0 ENERGY AND MINERAL RESOURCES	33
3.1 Introduction	33
3.2 Known Mineral and Energy Deposits	34
3.3 Known Mineral and Energy Prospects, Occurrences, and Mineralized Areas	37
3.4 Mining Claims, Leases and Material Sites	48
3.5 Mineral and Energy Deposit Types	52
3.6 Mineral and Energy Economics	62
4.0 LAND CLASSIFICATION FOR GEM RESOURCES POTENTIAL	64
4.1 Explanation of Classification Scheme	64
4.2 Classification of Appendicitus Hill (31-14) WSA	67
4.2.1 Locatable Resources	67
4.2.2 Leasable Resources	67
4.2.3 Saleable Resources	71
4.3 Classification of White Knob Mountains (31-17) WSA	71
4.3.1 Locatable Resources	71
4.3.2 Leasable Resources	73
4.3.3 Saleable Resources	74
4.4 Classification of Little Wood River (53-4) WSA	74
4.4.1 Locatable Resources	74
4.4.2 Leasable Resources	75
4.4.3 Saleable Resources	76
4.5 Classification of Friedman Creek (53-5) WSA	76
4.5.1 Locatable Resources	76
4.5.2 Leasable Resources	77
4.5.3 Saleable Resources	78






TABLE OF CONTENTS (Cont.)

	<u>Page</u>
5.0 RECOMMENDATIONS FOR FURTHER WORK	79
6.0 REFERENCES AND SELECTED BIBLIOGRAPHY	81
APPENDIX I: WILDERNESS STUDY AREA MAPS	90
APPENDIX II: FIELD VISIT REPORT	94
APPENDIX III: GEOCHEMICAL DATA	96




LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
1	Location Map	2
2	Topographic Map	4
3	Physiographic Setting	8
4	Geologic Map	10
5	Stratigraphic Column	11
6	Regional Structural Setting	23
7	Structural Units	24
8	Lineament Map	28
9	Regional Seismicity	29
10	Mineral Occurrences	35
11	Lava Creek District Geologic Map	39
12	Little Wood River District Geologic Map	43
13	Solid Muldoon Mine Geology	45
14	Geochemical Anomalies Firedman Creek WSA	50
15	Mining Claims Density Map	51
16	Oil and Gas Lease Status	53
17	Land Classification - Locatable Resources	68
18	Land Classification - Leasable Resources	69
19	Land Classification - Saleable Resources	72



LIST OF TABLES

<u>Table</u>		<u>Page</u>
I	Mineral Occurrences in the Pioneer Mountains GRA	36
II	Recorded Metal Production 1929-64, Little Wood River District	42
III	Anomalous Geochemical Samples, Friedman Creek WSA	49
IV	Oil and Gas Leases in WSA 31-14 and 31-17	54
V	Bureau of Land Management GEM Resources Classification System	65




PIONEER MOUNTAINS GRA1.0 INTRODUCTION

The Bureau of Land Management has adopted a two-phase procedure for the integration of geological, energy and minerals (GEM) resources data into the suitable/non-suitable decision-making process for Wilderness Study Area (WSAs). The objective of Phase I is the evaluation of existing data, both published and available unpublished data, for interpretation of the GEM resources potential of the WSAs. Wilderness Study Areas are grouped into areas based on geologic environment and mineral resources for initial evaluation. These areas are referred to as Geology, Energy, Mineral Resource Areas (GRAs).

The delineation of the GRAs is based on three criteria: (1) a 1:250,000 scale map of each GRA shall be no greater than 8½ x 11 inches; (2) a GRA boundary will not cut across a Wilderness Study Area; and (3) the geologic environment and mineral occurrences. The data for each GRA is collected, compiled, and evaluated and a report prepared for each GRA. Each WSA in the GRA is then classified according to GEM resources favorability. The classification system and report format are specified by the BLM to maintain continuity between regions.

This report is prepared for the Bureau of Land Management under contract number YA-553-CT2-1039. The contract covers GEM Region 2; Northern Rocky Mountains (Fig. 1). The Region includes 50 BLM Wilderness Study Areas



WGM Inc. Mining and Geological Consultants
Anchorage, Alaska

**BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS**

**Pioneer Mountains GRA,
Idaho
Location Map**

SCALE: 1"=45 MILES

DATA BY: WGM

DATE: 12/1982

DRAWN BY: DSI

APPROV: G.F.

FIGURE

1

• MOSCOW

• GRANDEVILLE

• MISSOULA

• ANACONDA

• BUTTE

• HELENA

• BOZEMAN

• DILLON

• SELMOY

**Pioneer Mountains
GRA**

• BOISE

• IDAHO FALLS

• POCAHELLO

TWIN FALLS RIVER

40 30 20 10 0 20 40 60 MILES
45 35 25 15 5 10 20 30 40

N E S W A D U T A N



totalling 583,182 acres. The WSAs were grouped into 22 GRAs for purposes of the Phase I GEM resources evaluation.

1.1 Location


The Pioneer Mountains GRA is located within 1s.2-6N., Rs.20-26E. (Fig. 2), Blaine and Butte Counties, central Idaho (Hailey and Idaho Falls NTMS Quadrangles). The GRA is within the Monument Resource Area of the Shoshone BLM district and the Big Butte Resource Area of the Idaho Falls BLM district. The GRA includes four BLM Wilderness Study Areas (WSAs): Appendicitus Hill (28,870 acres); White Knob Mountains (9,950 acres), Little Wood River (4,385 acres), and Friedman Creek (9,773 acres).

1.2 Population and Infrastructure



There are no towns within the Pioneer Mountains GRA. The nearest towns are Arco, five miles east of the GRA, and Hailey and Ketchum 8 and 12 miles west, respectively. State Highway 93 crosses the northeast corner of the GRA. Highway 20 passes 10 miles to the south of the GRA and Highway 75 lies to the west. There are several ranches and numerous gravel roads in the southern and eastern parts of the GRA.

1.3 Basis of the Report

This report is based on a compilation, review and analysis of the available published and unpublished data on the geology, energy and mineral resources of the Pioneer Mountains GRA. A number of geological and mineral resource



Approximate Boundary of Wilderness
Study Area.



**BLM Wilderness Study Areas Included in the
Pioneer Mountains GRA.**

<u>WSA</u>	<u>NAME</u>	<u>ACREAGE</u>
31-14	Appendicitis Hill	24,870
31-17	White Knob Mountains	9,950
53-4	Little Wood River	4,385
53-5	Friedman Creek	9,773

WGM Inc. Mining and Geological Consultants
Anchorage, Alaska

**BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS**

**Pioneer Mountains, Idaho
Topographic Map**

SCALE: 1"=4 MILES (1:250,000)

DATE BY WGM DATE 9/1982
DRAWN BY DBI APP'D G.F.

FIGURE

2




studies have been done in the area. Several of these were detailed studies which provide much useful data. BLM records of oil and gas leases and mining claims were reviewed and records of the Idaho Oil and Gas Commission were searched. WGM geologists spent two days in the field examining the Friedman Creek WSA during October 1982.

The data was compiled and reviewed by WGM project personnel and the Panel of Experts to produce the resource evaluation which comprises this report. Personnel involved in the report preparation are listed below.

Greg Fernette, Senior Geologist, WGM Inc.	Project Manager
C.G. Bigelow, President, WGM Inc.	Chairman, Panel of Experts
Joel Stratman, Geologist, WGM Inc.	Project Geologist
Jami Fernette, Land and Environmental Coordinator, WGM Inc.	Claims and Lease Compilation

Panel of Experts

C.G. Bigelow, President, WGM Inc.	Regional geology, metallic and minerals, mineral economics.
R.S. Fr�dericksen, Senior Geologist, WGM Inc.	Regional geology, metallic minerals.
David Blackwell, Ph.D., Professor Geophysics, Southern Methodist University	Geothermal.
Jason Bressler, Senior Geologist, WGM Inc.	Regional geology, metallic minerals.
Gary Webster, Ph.D., Chairman, Department of Geology, Washington State University	Oil and gas.
William Jones, Senior Geologist, WGM Inc.	Metallic minerals, coal, industrial minerals.
J.F. McOuat, President, Watts, Griffis & McOuat Ltd.	Mineral economics, and industrial minerals.




E.F. Evoy, Senior Geologist, Watts,
Griffis & McQuat Ltd.

Uranium and thorium.

1.4 Acknowledgements

We would like to thank Tim Carroll, District Geologist, Idaho Falls BLM for information on mining claims and mineral occurrences in the eastern part of the Pioneer Mountains GRA. Larry Dee loaned us copies of out-of-print publications and provided much assistance in the compilation phase of the project. We would like to thank Nance Petroleum, Billings, Montana, particularly Fred Brinkman, Vice President, and Keith Mohl, Geologist, for allowing Dr. Gary Webster to examine proprietary exploration data for the eastern part of the Pioneer Mountains GRA.




2.0 GEOLOGY


2.1 Introduction

The geology of the Pioneer Mountains GRA has been mapped by several U.S. Geological Survey workers (Dover, 1969; Nelson and Ross, 1969; U.S. Geological Survey and U.S. Bureau of Mines, 1981). Parts of the area are covered by University of Wisconsin M.S. theses (Bollmann, 1971; Lukowick, 1971; Volkmann, 1971; Wolbrink, 1970; Erwin, 1972). The geologic mapping has been compiled at a 1:250,000 scale by Rember and Bennett (1979a, b). These maps are the basis for Figure 4.

2.2 Physiography

The Pioneer Mountains GRA is in the northern Rocky Mountains physiographic province (Hunt, 1974) just north of the Snake River Plain (Fig. 3). The area includes parts of the Pioneer and White Knob Mountains. The topography is characterized by heavily dissected ridges and long structurally-controlled valleys. Drainage in the east half of the GRA is to the northeast via Antelope Creek. In the central part of the GRA drainage is to the south via Muldoon, Friedman and Fish Creeks. The valley floors have elevations of 5,800 to 6,000 feet above sea level. The highest elevation in the GRA is 11,505 feet on Smiley Mountain. The relief on most ridges is typically 1,800 to 2,500 feet.





Data from: Hunt (1974).

WGM Inc. Mining and Geological Consultants
Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS


Pioneer Mountains GRA,
Idaho
Physiographic Setting

SCALE: AS SHOWN	FIGURE
DATE: 3/1983	3
DRAWN BY: DSI	APPROVED: G.F.


2.3 Rock Units

The Pioneer Mountains GRA is in complexly thrust-faulted terrane; thus, the stratigraphy of the underlying rocks has only recently been partially unravelled (Skipp and Hait, 1977; Paull and Gruber, 1977; Dover, 1981). All of the Paleozoic (600-230 m.y.) section is allochthonous. The GRA encompasses parts of the four separate allochthons each of which is characterized by a different Paleozoic stratigraphic succession (Skipp and Hait, 1977); therefore, the Paleozoic rock units will be described for each allochthon. The four allochthons present in the GRA are (southwest to northeast) the: (1) Wood River, (2) Milligen, (3) Copper Basin, and (4) White Knob (Skipp and Hait, 1977). The stratigraphic column for these allochthons is illustrated in Figure 5.

The Wood River allochthon is composed entirely of the Wood River Formation (Fig. 5) which crops out in the west part of the Pioneer Mountains GRA and underlies about half of the Little Wood River WSA. The formation consists of a prominent basal conglomerate, known as the Hailey Conglomerate, about 370 feet thick (Hall et al., 1974). The conglomerate is composed of chert and quartzite pebbles and cobbles interbedded with brown limestone. The bulk of the formation is made up of over 3,000 feet of fine-grained, quartzose, limey sandstone to sandy limestone with quartzite interbeds, siltstone, and dark-gray siliceous argillite with minor easterly derived conglomerate (Hall et al., 1974; Skipp and Hall, 1975). The formation is distinguished by locally abundant fusulinid faunas (Bostwick, 1955; Hall et al., 1974) but also contains crinoid, bryozoan, brachiopod and algae fossils (Skipp et al., 1979). Based on its fossil content a Middle Pennsylvanian




Pioneer Mountains GRA, Idaho
Explanation for Figure 4



THE UNIVERSITY OF CHICAGO
LIBRARY





Source: Skipp and Halt, 1977
Skipp et. al., 1979

LEGEND

-  Not Present
-  Not Known

WGM Inc. Mining and Geological Consultants
Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS

**Pioneer Mountains GRA,
Idaho**
Stratigraphic Column

SCALE	DATE 9-83	FIGURE
DATA BY	ASVD	5
DRAWN BY	ASVD	

(305-290 m.y.) to Early Permian (280-251 m.y.) age is assigned to the Wood River (Skipp and Hait, 1977).

The Milligen allochthon is composed entirely of folded and faulted Milligen Formation (Fig. 5). The Milligen consists dominantly of slabby, phyllitic or subphyllitic argillite, and black, thin-bedded to massive, carbonaceous to siliceous, cherty argillite (Dover, 1981). Interbeds of gray, poorly sorted, fine-grained, quartzite, dolomitic siltstone and rare, thin beds of silty micritic to bioclastic limestone occur in the lower part of the unit. The Milligen Formation may be up to 4,000 feet thick, but the thickness is unknown due to structural disturbance. Sandberg et al. (1975) report an Early (395-370 m.y.) through early-Late Devonian (360-345 m.y.) age for the Milligen based on conodont faunas. The unit is interpreted to be a transitional, deep water, oceanic sequence deposited off the toe of a continental slope from a source area to the east (Skipp and Sandberg, 1975).

Like the previous two allochthons the Copper Basin allochthon is composed of a single thick unit, the Copper Basin Formation. The Copper Basin is structurally complicated; thus, its stratigraphic relationships are not completely understood. Dover (1981) divides the Copper Basin into three unnamed members: a lower clastic unit, a medial limestone, and an upper clastic unit.

The basal clastic unit and the medial limestone comprise the lower part of the Copper Basin Formation (Skipp and Hait, 1977). The lower clastic unit is at least 3,200 feet thick and consists of brown to green, medium- to thick-bedded, blocky argillite, and subordinate interbedded dark-gray

The first part of the report deals with the general situation of the country and the progress of the work during the year.

The second part of the report deals with the results of the work during the year.

The third part of the report deals with the financial situation of the country.

The fourth part of the report deals with the social situation of the country.

The fifth part of the report deals with the economic situation of the country.

The sixth part of the report deals with the political situation of the country.

The seventh part of the report deals with the cultural situation of the country.


The eighth part of the report deals with the military situation of the country.

The ninth part of the report deals with the foreign relations of the country.

The tenth part of the report deals with the future prospects of the country.

quartzite. The argillite is locally laminated and has graded bedding. The quartzite is poorly sorted with an argillaceous matrix and typically contains well rounded, coarse quartz grains or granules of black argillite floating in an argillaceous matrix. Conglomeratic beds with chert and siliceous argillite clasts occur locally (Dover, 1981). The middle limestone unit of the Copper Basin Formation as used by Dover (1981) is equivalent to the Drummond Mine Limestone, a key regional stratigraphic unit (Paull et al., 1972). The unit consists mainly of gray, yellow-brown weathering, slabby to platy, silty micritic limestone, which varies from silty limestone to calcareous siltstone. The rocks are locally silicified and recrystallized (Dover, 1981). The unit is about 2,650 feet thick (Paull et al., 1972). The lower contact of the limestone is gradational through 100 feet or more. Conodont, foraminifera, and ammonite fossils indicate an Early Mississippian (345-332 m.y.) age for the unit (Skipp and Bull, 1975). The Drummond Mine Limestone hosts much of the mineralization in the Copper Basin mining district (Dover, 1981). The lower part of the Copper Basin Formation comprises part of the Antler flysch (Poole, 1974).

The upper clastic unit or upper part of the Copper Basin Formation (Fig. 5) consists predominantly of argillite, gritty sandstone, and granule conglomerate. Graded bedding is common. Groove casts, rip-up conglomerate, cut and fill structures, and convoluted bedding indicate that the unit was deposited in an environment characterized by current scour, turbidity deposition, and sediment slumping. Fossils are scarce and poorly preserved, consequently determination of the age of the unit is tenuous. A Late Mississippian (332-310 m.y.) age is assigned based on stratigraphic relations with the underlying Drummond Mine Limestone and correlation with



more closely dated units in the region (Dover, 1981). The thickness of the upper part of the Copper Basin Formation is at least 2,500 feet. The Dry Fork thrust separates the lower and upper parts of the Copper Basin Formation.

The rocks within the White Knob allochthon are divided into a western and eastern facies. The western facies includes rocks ranging in age from Late Silurian (410-395 m.y.) to Pennsylvanian (310-280 m.y.). Rocks ranging in age from Devonian (395-345 m.y.) to Permian (280-230 m.y.) comprise the eastern facies (Fig. 5). The western facies is well exposed in the White Knob Mountains west of the Mackay and the eastern facies is recognized at Antelope Creek and Timbered Dome. The intervening interval between the two facies is covered by the Challis Volcanics; thus, it is possible that the two facies are separated by major faults (Skipp and Hait, 1977).

The oldest unit belonging to the eastern facies is the Late Silurian and Early Devonian Roberts Mountains Formation which is exposed in the Fish Creek Reservoir window (T.1N., R.22E.) about four miles south of the Pioneer Mountains GRA. The Roberts Mountains Formation consists of medium-gray, fossiliferous reef limestone, platy calcareous and dolomitic siltstone, and limestone conglomerate. Its thickness is estimated at about 65 feet (Skipp and Hall, 1975).

An Early to Late Devonian miogeosynclinal shelf carbonate sequence is also exposed in the Fish Creek Reservoir window. This sequence consists of (Fig. 5): (1) the Early and Middle (370-360 m.y.) Devonian Carey Dolomite, a light- to medium-gray, mostly clean, finely crystalline laminated dolomite

1. The first part of the report is a general introduction to the subject.

2. The second part is a detailed description of the methods used.

3. The third part is a discussion of the results obtained.

4. The fourth part is a conclusion and a summary of the findings.

5. The fifth part is a list of references.

6. The sixth part is a list of figures and tables.

7. The seventh part is a list of appendices.

8. The eighth part is a list of footnotes.

9. The ninth part is a list of acknowledgments.

10. The tenth part is a list of the author's address and contact information.

11. The eleventh part is a list of the author's previous work.

12. The twelfth part is a list of the author's future work.

13. The thirteenth part is a list of the author's publications.

14. The fourteenth part is a list of the author's awards and honors.

15. The fifteenth part is a list of the author's memberships in professional organizations.

16. The sixteenth part is a list of the author's other activities.

17. The seventeenth part is a list of the author's personal information.

18. The eighteenth part is a list of the author's family members.

19. The nineteenth part is a list of the author's pets.

20. The twentieth part is a list of the author's hobbies.

21. The twenty-first part is a list of the author's interests.

22. The twenty-second part is a list of the author's favorite books.

23. The twenty-third part is a list of the author's favorite movies.

24. The twenty-fourth part is a list of the author's favorite music.

25. The twenty-fifth part is a list of the author's favorite food.

26. The twenty-sixth part is a list of the author's favorite travel destinations.

27. The twenty-seventh part is a list of the author's favorite sports.

28. The twenty-eighth part is a list of the author's favorite games.

29. The twenty-ninth part is a list of the author's favorite TV shows.

30. The thirtieth part is a list of the author's favorite websites.

about 100 feet thick; (2) disconformably overlain by the Middle and Late Devonian Jefferson Dolomite, an 800 foot thick sequence of dolomite, dolomite conglomerate, and minor sandstone; (3) unconformably overlain by the Late Devonian Picabo Formation, a 190 foot thick sequence of gray fine-grained quartzose and calcareous sandstone, light-gray to pale-yellowish-brown sandy dolomite conglomerate, and minor gray fine-grained dolomite. The top of the Picabo is an unconformity (SkiPP and Hall, 1975).

The Cary Dolomite is also present in the Timber Dome area and is the oldest unit exposed in the Pioneer Mountains GRA. In the Timber Dome area, the Carey consists of about 330 feet of interbedded dolomite and quartzite. This area is within the eastern facies of the White Knob allochthon (SkiPP and Hait, 1977; Fig. 5).

Early Mississippian rocks unconformably overlie Late Devonian rocks throughout most of Idaho (SkiPP et al., 1979) including the Pioneer Mountains GRA. Mississippian rocks have been separated variously by different workers in the area surrounding the GRA. The following descriptions are based primarily on the regional division used by SkiPP et al. (1979). Many of these units have been recognized since completion of the detailed mapping of the MacKay Quadrangle (Nelson and Ross, 1969); thus, their distribution has not been mapped in detail throughout the GRA.

The Early and Late Mississippian McGowan Creek Formation is the basal Mississippian unit in the Pioneer Mountains GRA. The McGowan Creek correlates with the lower part of the Copper Basin Formation (SkiPP and Hait, 1977; Fig. 5). Regionally, the McGowan Creek consists of a lower

member of orogenic detritus, specifically fine-grained, thinly bedded turbidites, and an upper member of calcareous siltstone interbedded with silty micritic limestone, probably a starved-basin facies. The turbidite sequence ranges from about 100 feet thick in the Beaverhead Mountains to about 4,000 feet thick in the White Knob Mountains (Skipp et al., 1979). In the Timbered Dome area, the McGowan Creek is about 2,100 feet thick (Skipp and Hait, 1977). The upper member of the McGowan Creek ranges in thickness from 200 to almost 500 feet and is present only north and west of Hawley Mountain (about ten miles southeast of the Donkey Hills) (Skipp et al., 1979).

The McGowan Creek Formation is gradationally overlain by the White Knob Limestone. The White Knob is comprised of a lower sequence, about 1,150 feet thick, of interbedded silty micritic mudstone and calcareous siltstone which grades upward into thin- to thick-bedded, medium- to coarse-grained, fossiliferous bioclastic limestone. Eastward pointing tongues of conglomerate and sandstone related to the upper part of the Copper Basin Formation (Fig. 5) are interbedded with the bioclastic limestone in the upper 3,000 feet of the White Knob. Total thickness of the formation is about 5,450 feet (Skipp and Hait, 1977). The lower part of the White Knob represents forebank deposits whereas the upper part of the formation is the westernmost representative of a prograding carbonate bank. Locally abundant fossil evidence from the upper part of the White Knob indicates a Late Mississippian age for the unit (Skipp et al., 1979).

Carbonate bank and forebank deposits equivalent to the White Knob Limestone are divided into the Middle Canyon (oldest), Scott Peak, South Creek, and

Surrett Canyon Formations (youngest) from the White Knob Mountains east to the Montana-Idaho border. These formations represent a prograding carbonate-bank complex which generally thickens westward. Thickness ranges from 1,280 to 4,225 feet and is about 3,720 feet in the eastern part of the Pioneer Mountains GRA. The basal part of the complex, the Middle Canyon Formation, is a non-resistant calcareous sandy siltstone, and silty and sandy limestone weathering reddish brown. In the Pioneer Mountains GRA, the Middle Canyon is about 625 feet thick. It represents in part, the forebank deposit formed in front of the prograding bank; thus, it is correlative with the basal 1,150 feet of the White Knob Limestone. The succeeding Scott Peak Formation (1,700 feet thick in the GRA) and the uppermost Surrett Canyon Formation (1,000 feet thick in the GRA), both thick-bedded medium-dark-gray to dark-gray, variably cherty bioclastic limestones, represent a carbonate-bank accumulation. The intervening South-Creek Formation (395 feet thick in the GRA); a thin-bedded dark-gray micritic limestone probably indicates a period of relatively deep water marine circulation that temporarily interrupted carbonate-bank build-up. Fossil evidence indicates that all of the units belonging to the carbonate-bank complex are Late Mississippian in age (Skipp et al., 1979; Huh, 1967).

North of the Snake River Plain in east-central Idaho, about 4,600 feet of interbedded carbonate rocks, sandstone, siltstone and mudstone of latest Mississippian (345-310 m.y.) to Early Permian (280-251 m.y.) age are assigned to the Arco Hills (oldest), Bluebird Mountain, and Snaky Canyon (youngest) Formations. These rocks conformably overlie the Surrett Canyon Formation (Skipp et al., 1979). These rocks correspond to the unnamed carbonates shown in Figure 5.

The Arco Hills Formation is composed of interbedded medium-gray, olive-gray, yellowish-brown, and grayish-red thin-bedded, argillaceous, and silty or sandy limestone, calcareous mudstone or shale, siltstone and minor sandstone, and medium- to thick-bedded pure limestone. The proportion of shale and mudstone in the Arco Hills Formation increases from west to east. No shale or mudstone is present in the southern White Knob Mountains. The unit is a slope-former and is often buried beneath sandstone debris from the overlying Bluebird Mountain Formation. The upper contact with the Bluebird Mountain is sharp and is placed at the base of the stratigraphically lowest thick (12 feet or more) sandstone or siltstone bed. Brachiopods including Orbiculoidea sp., bryozoans, crinoid debris, and calcareous Foraminifera are abundant in the Arco Hills Formation. The Arco Hills is dated as Late Mississippian based on fossil evidence and stratigraphic position. In the southern White Knob Mountains the thickness of the formation is about 110 feet. The Arco Hills is not present in the western facies of the White Knob allochthon (Skipp et al., 1979).

In the Pioneer and White Knob Mountains, the Bluebird Mountain Formation consists of calcareous siltstone, very fine-grained sandstone, and interbedded sandy limestone. The upper contact with the overlying Snaky Canyon Formation is gradational and placed at the position in the section above which limestones are predominate. Foraminifera and the conodonts, Gnathodus bilineatus and Cavusgnathus sp., indicate that the Bluebird Mountain is Late Mississippian. The Bluebird Mountain is 25 feet thick in the southern White Knob Mountains. The thickness the Bluebird in the western facies is probably somewhat less (Skipp et al., 1979).

The first part of the report deals with the general situation of the country. It is a very interesting and detailed account of the country's history and its present state. The author has done a great deal of research and has gathered a wealth of material. The second part of the report deals with the political situation. It is a very interesting and detailed account of the country's political system and its present state. The author has done a great deal of research and has gathered a wealth of material. The third part of the report deals with the economic situation. It is a very interesting and detailed account of the country's economic system and its present state. The author has done a great deal of research and has gathered a wealth of material. The fourth part of the report deals with the social situation. It is a very interesting and detailed account of the country's social system and its present state. The author has done a great deal of research and has gathered a wealth of material. The fifth part of the report deals with the cultural situation. It is a very interesting and detailed account of the country's cultural system and its present state. The author has done a great deal of research and has gathered a wealth of material. The sixth part of the report deals with the military situation. It is a very interesting and detailed account of the country's military system and its present state. The author has done a great deal of research and has gathered a wealth of material. The seventh part of the report deals with the foreign relations situation. It is a very interesting and detailed account of the country's foreign relations system and its present state. The author has done a great deal of research and has gathered a wealth of material. The eighth part of the report deals with the future of the country. It is a very interesting and detailed account of the country's future and its present state. The author has done a great deal of research and has gathered a wealth of material.

The report is a very interesting and detailed account of the country's history and its present state. The author has done a great deal of research and has gathered a wealth of material. The report is a very interesting and detailed account of the country's political system and its present state. The author has done a great deal of research and has gathered a wealth of material. The report is a very interesting and detailed account of the country's economic system and its present state. The author has done a great deal of research and has gathered a wealth of material. The report is a very interesting and detailed account of the country's social system and its present state. The author has done a great deal of research and has gathered a wealth of material. The report is a very interesting and detailed account of the country's cultural system and its present state. The author has done a great deal of research and has gathered a wealth of material. The report is a very interesting and detailed account of the country's military system and its present state. The author has done a great deal of research and has gathered a wealth of material. The report is a very interesting and detailed account of the country's foreign relations system and its present state. The author has done a great deal of research and has gathered a wealth of material. The report is a very interesting and detailed account of the country's future and its present state. The author has done a great deal of research and has gathered a wealth of material.

The Mississippian-Pennsylvanian contact is gradational in the Pioneer Mountains GRA, and the Pennsylvanian rocks are represented by the lower part of the recently defined Snaky Canyon Formation (Skipt et al., 1979) which is comprised of the basal Bloom Member, the succeeding Gallagher Peak Sandstone Member, and the lower part of the overlying Juniper Gulch Member. Only the basal Bloom Member has been recognized in the White Knob allochthon. An incomplete faulted section of the Snaky Canyon at Timbered Dome is 2,855 feet thick (Skipt et al., 1979). Skipt and Hait (1977) report a thickness of 330+ feet for the western facies of the Snaky Canyon. The Bloom Member consists of medium-gray limestone, largely sandy or silty, interbedded with thin beds of very fine-grained, yellowish-brown weathering sandstone and siltstone. Medium- to coarse-grained chert and quartz sandstone, shaly mudstone and siltstone, as well as minor conglomerate are also present in the Bloom Member in the southern White Knob Mountains. Fossil evidence indicates that the Bloom Member spans Early (310-305 m.y.) and Middle Pennsylvanian (305-290 m.y.) time (Skipt et al., 1979).

In the southern White Knob Mountains, along Antelope Creek, at least 164 feet of dark-gray thin-bedded argillaceous and silty limestone, some of it phosphatic, is present in a small graben. The fauna consisting of brachiopods, bryozoans, and mollusks indicate an age (Early Permian) equivalent to the Meade Peak Member of the Phosphoria Formation (Skipt et al., 1979). The contact with the underlying Snaky Canyon is not exposed, but it is presumably conformable.

Mesozoic (240-65 m.y.) clastic rocks rest unconformably on pre-Mesozoic rocks in the Pioneer Mountains GRA. These rocks are mostly pebble- to

The first part of the report deals with the general situation of the country. It is a very interesting and informative study of the country's development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's development. It is a must-read for anyone interested in the country's development.

The second part of the report deals with the economic situation of the country. It is a very interesting and informative study of the country's economic development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's economic development. It is a must-read for anyone interested in the country's economic development.

The third part of the report deals with the social situation of the country. It is a very interesting and informative study of the country's social development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's social development. It is a must-read for anyone interested in the country's social development.

The fourth part of the report deals with the political situation of the country. It is a very interesting and informative study of the country's political development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's political development. It is a must-read for anyone interested in the country's political development.

The fifth part of the report deals with the cultural situation of the country. It is a very interesting and informative study of the country's cultural development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's cultural development. It is a must-read for anyone interested in the country's cultural development.

The sixth part of the report deals with the environmental situation of the country. It is a very interesting and informative study of the country's environmental development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's environmental development. It is a must-read for anyone interested in the country's environmental development.

The seventh part of the report deals with the international situation of the country. It is a very interesting and informative study of the country's international development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's international development. It is a must-read for anyone interested in the country's international development.

The eighth part of the report deals with the future of the country. It is a very interesting and informative study of the country's future development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country's future development. It is a must-read for anyone interested in the country's future development.

boulder-conglomerates composed of well-rounded clasts representing stream gravels and conglomerate or breccia consisting of angular fragments representing talus or other colluvial deposits. The most abundant rock type in these clastic deposits is comprised of quartzite clasts set in a poorly sorted, moderately well cemented quartz-sand matrix. Plant fossils found in the conglomerates indicate a Late Cretaceous (100-65 m.y.) to Paleocene (65-55 m.y.) age for these sediments (Dover, 1981).

The Eocene (55-38 m.y.) Challis Volcanics unconformably overlie the Mesozoic clastics and the Paleozoic rocks in the Pioneer Mountains GRA. The Challis Volcanics are predominantly intermediate flows and tuffaceous rocks (Skipp, 1981). The bulk of the volcanic rocks consist of dark colored porphyritic flows, tuffs and volcanic breccia (Nelson and Ross, 1969). The flows are gray, brown, purple and green hornblende andesites and latites (Dover, 1981). The majority of the Challis Volcanics were extruded between 51 and 45 m.y. ago based on potassium-argon dates (Skipp, 1981).

The layered rocks of the Pioneer Mountains GRA are intruded by two stocks related to the Challis episode of volcanism: (1) the Lake Creek stock and (2) the Muldoon stock (Dover, 1981). The Lake Creek stock is an elongate, northwest-trending body exposed in the north-central part of the GRA (Nelson and Ross, 1969; Fig. 4). The stock is composed of pyroxene-hornblende-biotite quartz monzonite. Phenocrysts of plagioclase, pyroxene and biotite occur in a fine-grained quartz-potassium feldspar matrix. The rock is generally altered. Hornblende is biotitized and biotite is chloritized. The wall rocks are often bleached, silicified and hornfelsed at contacts with the intrusive (Dover, 1981). Armstrong (1975) reports a radiometric

1. The first part of the report deals with the general situation of the country and the progress of the work during the year. It is divided into two main sections: the first section deals with the general situation and the second section deals with the progress of the work.

2. The second part of the report deals with the results of the work during the year. It is divided into two main sections: the first section deals with the results of the work in the field of research and the second section deals with the results of the work in the field of education.

3. The third part of the report deals with the conclusions of the work during the year. It is divided into two main sections: the first section deals with the conclusions of the work in the field of research and the second section deals with the conclusions of the work in the field of education.

4. The fourth part of the report deals with the recommendations of the work during the year. It is divided into two main sections: the first section deals with the recommendations of the work in the field of research and the second section deals with the recommendations of the work in the field of education.


5. The fifth part of the report deals with the summary of the work during the year. It is divided into two main sections: the first section deals with the summary of the work in the field of research and the second section deals with the summary of the work in the field of education.

lead age of 50 to 48 m.y. for the stock. Dover (1981) describes contact relations which suggest that the stock is slightly younger than the Challis Volcanics. The Muldoon stock, exposed in two places above Muldoon Creek, is also an elongate northwest-trending intrusive (Fig. 4). It is similar in composition to the Lake Creek stock but has a medium-grained groundmass. Biotite from the Muldoon stock has been dated as 47 m.y. old by potassium-argon methods (Dover, 1981). Small bodies of quartz monzonite present in the Lava Creek mining district may indicate the presence of another near-surface stock (Anderson, 1929).

Rhyolite and andesite dikes occur locally in the Pioneer Mountains GRA (Nelson and Ross, 1969; Dover, 1981). The dikes occur in fracture zones especially those trending northeast and east. Some of the dikes were feeders for the Challis Volcanics.

Pleistocene (2-0.1 m.y.) basalt flows which are part of the Snake River Group occur in two areas located on the southern margin of the Pioneer Mountains GRA. These rocks are dark-gray to black, tube-fed pahoehoe lava flows, and bedded moderately oxidized scoria, cinders, and ash (Kuntz, 1978). These flows were fed from a vent located about four miles west of Arco. The lavas rest on and are partly buried by alluvial deposits of late Pleistocene age.


Pleistocene alluvium, glacial and lake deposits overlie all of the rocks in the area. The Pleistocene deposits are dominantly glacial and glacio-fluvial in origin (Dover, 1981). The youngest deposits in the area are Holocene (0.1 m.y. to present) alluvial sediments.




2.4 Structural Geology and Tectonics

Regionally the Pioneer Mountains GRA is within the frontal Sevier orogenic belt of the Cordilleran fold-thrust belt (Fig. 6). In the area northwest of the Snake River Plain and east of the Idaho batholith, a minimum of nine major allochthons are recognized including (from west to east): the Wood River, Milligen, Copper Basin, White Knob, Lost River-Arco Hills, Lemhi, Beaverhead, Medicine Lodge and Tendoy allochthons. Each of these allochthons has an internally consistent stratigraphy that differs from adjacent plates. All of the allochthons, except the structurally highest Wood River, are bounded above and below by thrust faults. The allochthons can be grouped into three stacks consisting of: (1) the Pioneer or western stack, structurally the highest, comprising the Wood River, Milligen, Copper Basin and White Knob allochthons, (2) the Lemhi or middle stack comprising the Lost River-Arco Hills, Lemhi, and Beaverhead allochthons, and (3) the Lima or eastern stack, structurally the lowest, comprising the Medicine Lodge and Tendoy allochthons (Skiipp and Hait, 1977). The Pioneer Mountains GRA encompasses parts of all the allochthons belonging to the Pioneer stack (Fig. 7).

The Wood River allochthon occupies an area 50 miles long in a north-northwest trending direction. Its western boundary is marked by the eastern edge of the Idaho batholith, and it is about 25 miles wide in the Wood River area. The allochthon is about 3,000 feet thick and is composed entirely of the Wood River Formation (Fig. 5). It has a minimum stratigraphic separation of 9,800 feet with respect to the Milligen Formation in the underlying Milligen allochthon. Facies differences and stratigraphic separation indicate that the Wood River allochthon was transported a minimum distance






Data From:

Skip and Hail, (1977)

Dover, (1981)



LEGEND



Alluvium



Challis Volcanics



Intrusive Rocks

MWR - Milligan/Wood River Allochthon

CB - Copper Basin Allochthon

(CBP) Copper Basin Plate

(GMP) Glide Mountain Plate

WK - White Knob Allochthon

WGM Inc. Mining and Geological Consultants
Anchorage, Alaska


BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS

Pioneer Mountains, Idaho

Structural Units

SCALE: _____
DATE BY: DATE 9-83 NAME _____
DRAWN BY: ADVICE: _____

FIGURE
7



of 10 miles on the Wood River thrust. The Wood River allochthon is always associated with the underlying Milligen allochthon; therefore, the two may have moved as a unit during the most recent thrusting (Skipp and Hall, 1975). The Wood River allochthon is deformed and contact metamorphosed by the Late Cretaceous Idaho batholith; thus, latest eastward movement was after Early Permian time and pre-Late Cretaceous. Movement probably occurred during the Sevier orogeny (Hall et al., 1975).

The Milligen allochthon occupies about the same geographic area as the Wood River allochthon. The Milligen allochthon is about 4,000 feet thick and consists exclusively of tightly folded and faulted Milligen Formation. The eastern edge of the allochthon is tightly folded and overturned. The Milligen allochthon is likely thrust over the western part of the Copper Basin allochthon and the lower Paleozoic miogeoclinal carbonates in the Fish Creek Reservoir area. The juxtaposition of Devonian continental rise and miogeoclinal sequences indicates a minimum eastward transport of 30 miles (Skipp and Hait, 1977). If the continental margin occupied a position near the western margin of the Idaho batholith as suggested by Armstrong et al. (1977), then transport of at least 80 miles has occurred (Skipp and Hait, 1977). Regional studies of the Milligen Formation indicate that the unit was deformed and thrust eastward during the Antler orogeny in Late Devonian and Early Mississippian time (Hall et al., 1975). Like the overlying Wood River allochthon, the Milligen allochthon is deformed and thermally metamorphosed by the Idaho batholith; therefore, most recent movement is the same as for the Wood River allochthon, probably Sevier (Skipp and Hait, 1977).

The first of these is the fact that the
the second is the fact that the
the third is the fact that the
the fourth is the fact that the
the fifth is the fact that the
the sixth is the fact that the
the seventh is the fact that the
the eighth is the fact that the
the ninth is the fact that the
the tenth is the fact that the
the eleventh is the fact that the
the twelfth is the fact that the
the thirteenth is the fact that the
the fourteenth is the fact that the
the fifteenth is the fact that the
the sixteenth is the fact that the
the seventeenth is the fact that the
the eighteenth is the fact that the
the nineteenth is the fact that the
the twentieth is the fact that the
the twenty-first is the fact that the
the twenty-second is the fact that the
the twenty-third is the fact that the
the twenty-fourth is the fact that the
the twenty-fifth is the fact that the
the twenty-sixth is the fact that the
the twenty-seventh is the fact that the
the twenty-eighth is the fact that the
the twenty-ninth is the fact that the
the thirtieth is the fact that the
the thirty-first is the fact that the
the thirty-second is the fact that the
the thirty-third is the fact that the
the thirty-fourth is the fact that the
the thirty-fifth is the fact that the
the thirty-sixth is the fact that the
the thirty-seventh is the fact that the
the thirty-eighth is the fact that the
the thirty-ninth is the fact that the
the fortieth is the fact that the
the forty-first is the fact that the
the forty-second is the fact that the
the forty-third is the fact that the
the forty-fourth is the fact that the
the forty-fifth is the fact that the
the forty-sixth is the fact that the
the forty-seventh is the fact that the
the forty-eighth is the fact that the
the forty-ninth is the fact that the
the fiftieth is the fact that the
the fifty-first is the fact that the
the fifty-second is the fact that the
the fifty-third is the fact that the
the fifty-fourth is the fact that the
the fifty-fifth is the fact that the
the fifty-sixth is the fact that the
the fifty-seventh is the fact that the
the fifty-eighth is the fact that the
the fifty-ninth is the fact that the
the sixtieth is the fact that the
the sixty-first is the fact that the
the sixty-second is the fact that the
the sixty-third is the fact that the
the sixty-fourth is the fact that the
the sixty-fifth is the fact that the
the sixty-sixth is the fact that the
the sixty-seventh is the fact that the
the sixty-eighth is the fact that the
the sixty-ninth is the fact that the
the seventieth is the fact that the
the seventy-first is the fact that the
the seventy-second is the fact that the
the seventy-third is the fact that the
the seventy-fourth is the fact that the
the seventy-fifth is the fact that the
the seventy-sixth is the fact that the
the seventy-seventh is the fact that the
the seventy-eighth is the fact that the
the seventy-ninth is the fact that the
the eightieth is the fact that the
the eighty-first is the fact that the
the eighty-second is the fact that the
the eighty-third is the fact that the
the eighty-fourth is the fact that the
the eighty-fifth is the fact that the
the eighty-sixth is the fact that the
the eighty-seventh is the fact that the
the eighty-eighth is the fact that the
the eighty-ninth is the fact that the
the ninetieth is the fact that the
the ninety-first is the fact that the
the ninety-second is the fact that the
the ninety-third is the fact that the
the ninety-fourth is the fact that the
the ninety-fifth is the fact that the
the ninety-sixth is the fact that the
the ninety-seventh is the fact that the
the ninety-eighth is the fact that the
the ninety-ninth is the fact that the
the hundredth is the fact that the

The Copper Basin allochthon extends from Fish Creek Reservoir on the southwest to the eastern flank of Blizzard Mountain (just north of Craters-of-the-Moon) on the southeast. The allochthon is about 12 miles wide and its thickness ranges from 0 just west of Fish Creek Reservoir to 4,500 feet in the southern Pioneer Mountains. The Copper Basin allochthon is imbricated and displaced by N70°E trending tear faults. These faults have left-lateral offset. The allochthon is also cut by normal faults having relative movements of less than 1,000 feet. The latter faults may be caused by Tertiary uplift associated with diapiric structures in the central and southern Pioneer Mountains. The structurally high Glide Mountain plate (Fig. 7) is included in the Copper Basin allochthon. The relatively flat Glide Mountain thrust may override the steeper Muldoon Canyon fault. The sole fault at the base of the Copper Basin allochthon, the Fish Creek-Copper Basin thrust has overridden Late Silurian to Mississippian rocks belonging to the White Knob allochthon. The movement on the Copper Basin allochthon is difficult to estimate, but the complex imbricate thrusting, the great thickness of the Copper Basin Formation, and the tectonically reversed stratigraphic order within the allochthon suggest a minimum eastward transport of 12 miles. On the west, the Copper Basin allochthon is overridden by the Wood River and Milligen allochthons which were in place by pre-Late Cretaceous time (Skipp and Hait, 1977).

The White Knob allochthon extends from the trace of the Copper Basin thrust on the west to the Big Lost River Valley on the east. The basal thrust of the allochthon is not exposed in the Big Lost River Valley, but the fault is needed to explain the facies differences between rocks present in the southern White Knob Mountains and those present in the southern Lost River

The first of these is the fact that the data are not normally distributed. The second is that the data are not independent. The third is that the data are not stationary.

The first of these is the fact that the data are not normally distributed. The second is that the data are not independent. The third is that the data are not stationary.

The first of these is the fact that the data are not normally distributed. The second is that the data are not independent. The third is that the data are not stationary.

The first of these is the fact that the data are not normally distributed. The second is that the data are not independent. The third is that the data are not stationary.

The first of these is the fact that the data are not normally distributed. The second is that the data are not independent. The third is that the data are not stationary.

The first of these is the fact that the data are not normally distributed. The second is that the data are not independent. The third is that the data are not stationary.

The first of these is the fact that the data are not normally distributed. The second is that the data are not independent. The third is that the data are not stationary.

The first of these is the fact that the data are not normally distributed. The second is that the data are not independent. The third is that the data are not stationary.

The first of these is the fact that the data are not normally distributed. The second is that the data are not independent. The third is that the data are not stationary.

The first of these is the fact that the data are not normally distributed. The second is that the data are not independent. The third is that the data are not stationary.


Range. At the surface, the allochthon is 15 to 20 miles wide. The thickness is not known with certainty, but Skipp and Hait (1977) estimate it at 10,000 feet. At the mouth of Antelope Creek, crustal shortening of 40% is estimated for the tight-folding present; thus, a minimum distance of at least 5 miles of eastward transport is inferred. However, an exotic block of granitic gneiss in the Challis Volcanics near Antelope Creek and the presence of high grade Precambrian(?) metamorphic rocks in the structural position of the White Knob allochthon in the central Pioneer Mountains, both suggest that eastward transport may have been much greater. The White Knob allochthon is broken by many north-south and northwest-southeast trending normal faults of basin-and-range affinities (Skipp and Hait, 1977).

Most of the major faults in the Pioneer Mountains GRA form lineaments visible on LANDSAT photos (Fig. 8). Two sets of circular features, one on Dry Fork Creek in the south-central part of the GRA and the other southeast of WSA 53-4, may indicate the presence of near-surface intrusives. Circular fault patterns are associated with buried stocks northeast of the GRA (Ruppel, 1982).

Geologic hazards in the Pioneer Mountains GRA relate mainly to seismicity. The area is just south of the Sawtooth Mountains earthquake swarm (Fig. 9) within the Intermountain Seismic belt (Smith and Sbar, 1974).



2.5 Paleontology

Fossils are reported in many of the sedimentary units in the Pioneer Mountains GRA but no special collecting or index localities are reported.





Approximate Boundary of Wilderness
Study Area.



Legend

— LINEAMENT

Data from: Ruppel (1982), Mitchell et al., (1980),
Day (1974), Haskett (1974).


WGM Inc. Mining and Geological Consultants
Anchorage, Alaska


BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS

Pioneer Mountains, Idaho

Lineament Map

SCALE: 1" = 4 MILES (1:250,000)				FIGURE 8
DATE BY	DATE	9/1982	REVIEW	
DRAWN BY	DSI	APPROV.	G.F.	





EPICENTERS FOR INTERMOUNTAIN SEISMIC BELT SHOWING ALL EARTHQUAKES REPORTED BY NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION (NOAA) OF ABOUT MAGNITUDE 3.0 OR GREATER FROM 1961 THROUGH 1970. DOUBLE ARROW SHOWS ZONE OF LOW SEISMICITY. CIRCLES MARK EARTHQUAKE SWARMS: 1, FLATHEAD LAKE, MONT.; 2, HELENA, MONT.; 3, YELLOWSTONE PARK; 4, SAWTOOTH RANGE, IDAHO; 5, IDAHO BATHOLITH; 6, CARIBOU RANGE, IDAHO; 7, GREAT SALT LAKE DESERT, UTAH; 8, LEHMAN CAVE MONUMENT, NEVADA; 9, CEDAR CITY, UTAH.

DATA FROM: SMITH AND SBAR (1974)


0 70 140 210 MILES
SCALE APPROXIMATE

WGM Inc. Mining and Geological Consultants
Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS

**Pioneer Mountains GRA,
Idaho
Regional Seismicity**

SCALE	DATE: 3/1983	REVISION	FIGURE
DATA BY: WGM	APPROV: G.F.		9
DRWN BY: DSI			



Conodonts and foraminifera in the Drummond Mine Limestone are significant in determining the age of the Copper Basin Formation (Dover, 1981). Foraminifera, brachiopods, bryozoans and corals are common in the White Knob Limestone and are described in detail by Skipp et al. (1979). Fossil fragments, crinoid columnals, bryozoans, and brachiopods are common in parts of the conglomeratic lower Wood River Formation (Thomasson, 1959). Fusulinids are abundant in some units of the upper part of the Wood River Formation (Hall et al., 1974). Plant fragments are found in the conglomerate at the base of the Challis Volcanics (Paull, 1974).

2.6 Historical Geology

Beginning in Middle Devonian a deep basin developed in central Idaho. To the east a miogeosyncline developed which eventually became distinctly differentiated from the semi-stable cratonic shelf along a sharp hinge line near the present Lemhi Range. Seas were regionally transgressive during all of Middle and part of Late Devonian time (Scholten and Hait, 1962).

Latest Devonian to Early Mississippian was a period of upheaval associated with the Antler orogeny. Devonian and older rocks were uplifted and eroded as the Antler highland developed in southwestern Idaho and northern Nevada (Skipp, 1981). Some thrust faulting accompanied the Antler orogeny as indicated by the early movement of the Milligen allochthon (Skipp and Hait, 1977). Turbidites filled a deep and narrow flysch trough developed adjacent to the eastern margin of the Antler highland. East of the flysch trough, thick orogenic, thin starved basin, and thick carbonate sequences were

1. The first part of the report deals with the general situation of the country and the position of the various groups.

2. The second part of the report deals with the economic situation and the position of the various groups.

3. The third part of the report deals with the social situation and the position of the various groups.

4. The fourth part of the report deals with the political situation and the position of the various groups.

5. The fifth part of the report deals with the cultural situation and the position of the various groups.

6. The sixth part of the report deals with the religious situation and the position of the various groups.

7. The seventh part of the report deals with the legal situation and the position of the various groups.

8. The eighth part of the report deals with the administrative situation and the position of the various groups.

9. The ninth part of the report deals with the judicial situation and the position of the various groups.

10. The tenth part of the report deals with the military situation and the position of the various groups.

11. The eleventh part of the report deals with the foreign relations situation and the position of the various groups.

12. The twelfth part of the report deals with the internal security situation and the position of the various groups.

13. The thirteenth part of the report deals with the public health situation and the position of the various groups.

14. The fourteenth part of the report deals with the education situation and the position of the various groups.


15. The fifteenth part of the report deals with the labor situation and the position of the various groups.

deposited in an outer cratonic platform or foreland basinal environment. Sedimentation continued without major interruption through Mississippian time (Skipp et al., 1979). The flysch basin had filled by Early Pennsylvanian time and was succeeded by a carbonate bank. Coincident with the rise of the ancestral Rocky Mountains during Middle Pennsylvanian time, the sediments in the western portion of the flysch basin were uplifted to form the Copper Basin highland. Minor coarse detritus from this highland was shed eastward into the shallow water carbonate bank regime of the Snake Canyon Formation (Skipp, 1981).


The time gap between Permian rocks and the Eocene Challis Volcanics represents the Sevier orogeny. During Early Cretaceous (141-100 m.y.) to Paleocene the major thrust sheets developed forming the allochthonous terrane presently underlying the Pioneer Mountains GRA (Skipp, 1981). Eastward telescoping due to thrusting across the northeast margin of the Snake River Plain is estimated to have produced crustal shortening in excess of 180 miles, and possibly as much as 400 miles. Major thrust movement, at least in the Pioneer Mountains had ceased by Late Cretaceous time when the Idaho batholith was intruded (Skipp and Hait, 1977).

The major part of the Eocene Challis Volcanics was erupted from about 51 to 45 m.y. ago. Minor volcanism and associated intrusive activity continued in the south-central Idaho region until about 40 m.y. ago. Normal and strike-slip faulting proceeded and accompanied volcanism (Skipp, 1981).

After a tectonically relatively inactive period during Oligocene (38-23 m.y.) and early Miocene (23-6 m.y.) time, major basin-and-range extension




faulting commenced about 17 m.y. ago. Eastward tilting of the uplifted fault blocks along moderately dipping normal faults located on their west flank formed the narrow, linear north- to northwest-trending mountain ranges (Skipp, 1981). The inception of basin-and-range faulting was followed closely by the initial downwarping and faulting of the north margin of the Snake River Plain, located south of the Pioneer Mountains GRA (Pankrantz and Ackerman, 1982). Bimodal volcanism related to the development of the Snake River Plain began at least 14 m.y. ago (Christiansen and McKee, 1978) and continued to less than 20,000 years ago (Kuntz et al., 1979).



3.0 MINERAL AND ENERGY RESOURCES

3.1 Introduction

Information on the mineral and energy resources of the Pioneer Mountains GRA was collected from all available sources. General mineral resource data was obtained from the U.S. Geological Survey CRIB File, the U.S. Bureau of Mines MILS File, and the Idaho Bureau of Mines Mines and Prospects Maps Series (Strowd et al., 1981; Hustedde et al., 1981). More specific mineral resource information is available in reports by Anderson (1929), Anderson and Wagner (1946), Finch (1917), Umpleby (1917), and Umpleby et al. (1930). The U.S. Geological Survey and the U.S. Bureau of Mines recently completed an evaluation of the Boulder-Pioneer Wilderness Study Area, which includes the northern part of the GRA (U.S. Geological Survey and U.S. Bureau of Mines, 1981). This report was of great value in this study. Notes on mineral occurrences in the eastern part of the GRA by Tim Carroll also were valuable. Energy resource data was obtained from Breckenridge (1982), the Idaho Oil and Gas Commission, and Mitchell et al. (1980). Nance Petroleum of Billings, Montana kindly allowed WGM to examine data pertaining to the eastern part of the GRA. Two days of field work were completed in the vicinity of the Friedman Creek WSA in October 1982. This work included the collection and analysis of 33 stream sediment samples from streams draining the Friedman Creek WSA.




3.2 Known Mineral and Energy Deposits

There are over 70 mineral deposits, prospects, and occurrences in the Pioneer Mountains GRA (Fig. 10, Table I). Most of the deposits are in the Lava Creek, Copper Basin, and Little Wood River mining districts.

The Lava Creek district (Fig. 11), located in the south-central part of the Pioneer Mountains GRA, is described by Anderson (1929). The mines in the Lava Creek district produced about \$500,000 worth of silver and lead during 1884-1886. Within the GRA, the district encompasses two groups of deposits; an eastern group on Champagne Creek (loc. 6-12, Fig. 10, Table I) and a western group on Leadbelt and Dry Fork Creeks (loc. 13-14, Fig. 10). About half of the production from the district came from the Hornsilver mine (loc. 4, Fig. 11). The Lead Belt mine (loc. 13, Fig. 10, Table I) has also produced a small amount of silver and lead (Simons, 1981). There is little data on the grade of the ore. Anderson (1929) reports that the ore at the Last Chance mine (loc. 8, Fig. 11) averaged 125 oz/ton silver and 4% copper. Most of the ore in the district was mined from oxidized silver-rich secondary deposits near the surface (Anderson, 1929).


The lodes occur in fissure veins or brecciated zones in volcanics or limestone. Replacement of breccia fragments and wall rocks along vein walls is common. Mineralization consists of argentiferous galena, sphalerite, pyrite, and minor chalcopyrite in a gangue consisting of quartz and locally calcite. The veins range from a stringer up to 12 feet in width. The main vein at the Hornsilver mine averaged four feet in width. Alteration often



Ag - Silver
 Au - Gold
 Bi - Bismuth
 Cu - Copper
 Pb - Lead
 Sb - Antimony
 W - Tungsten
 Zn - Zinc
 g - Sand and gravel




Approximate Boundary of Wilderness Study Area.



LEGEND

- Mineral Orebody
- Mineral Deposit
- Mineral Occurrence
- ✕ Active Quarry
- ✕ Inactive Quarry
- ✕ Active gravel or clay pit
- ✕ Inactive gravel or clay pit

1 occurrence number
 Cu commodity



WGM Inc. Mining and Geological Consultants
 Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT

REGION 2 NORTHERN ROCKY MOUNTAINS

Pioneer Mountains, Idaho

Mineral Occurrences


SCALE 1" = 4 MILES	FIGURE
DATA BY WGM DATE 1/1983	10
DRAWN BY DBT	APPROV. G.F.




TABLE I

MINERAL OCCURRENCES IN THE PIONEER MOUNTAINS GRA, IDAHO


Map No.	Name	Commodity	Source of Data
1	Mel Johnson placer	Agate	Carroll, 1981
2	Unnamed	?	Carroll, 1981
3	Unnamed	?	Carroll, 1981
4	Unnamed	?	Carroll, 1981
5	Espinoza claims	Copper minerals and Agate	Carroll, 1981
6	Ella mine	Pb-Zn-Ag-Cu-Au	Anderson, 1929
7	Hornsilver mine	Pb-Zn-Ag-Cu-Au + Sb-Bi	Anderson, 1929
8	Valley Arco mine	?	Anderson, 1929
9	Reliance mine	Pb-Zn-Bi-Cu-Ag	Anderson, 1929
10	St. Louis mine	Pb-Zn-Bi-Cu-Ag	Anderson, 1929
11	Oxide Lode prospect	Fe	Anderson, 1929
12	Last Chance mine	Zn-Ag-Cu	Anderson, 1929
13	Lead Belt mine	Pb-Ag-Au	Anderson, 1929
14	Butte-Antelope mine	Pb-Ag-Au-Cu	Anderson, 1929
15	Unnamed	?	Strowd et al., 1981
16	Unnamed	?	Strowd et al., 1981
17	Unnamed	?	Strowd et al., 1981
18	Unnamed	?	Strowd et al., 1981
19	Unnamed	?	Strowd et al., 1981
20	Unnamed	?	Strowd et al., 1981
21	Upper Lake Creek	W-Ag	Strowd et al., 1981
22	Unnamed	Ag-Pb	Tuchek and Ridenour, 1981
23	Unnamed	Ag-Pb	Tuchek and Ridenour, 1981
24	Cornhusker group	Pb-Zn-Cu	Strowd et al., 1981
25	Candy Cane group	Pb-Zn-Cu	Tuchek and Ridenour, 1981
26	Silver Bell	Pb-Ag-Cu	Tuchek and Ridenour, 1981
27	Unnamed	Ag-Cu-Pb	Tuchek and Ridenour, 1981
28	Unnamed	Barite	Tuchek and Ridenour, 1981
29	Star Hope mine	Pb-Zn-Ag-Cu-Au	Tuchek and Ridenour, 1981
30	Unnamed	?	Strowd et al., 1981
31	Bent Pine Tree group	Ag-Pb-Zn-Cu	Tuchek and Ridenour, 1981
32	Drummond mine	Cu-Au-Pb-Ag	Umpleby, 1917
33	Silver Mint Nos. 1 and 2	Cu-Au-Pb	Tuchek and Ridenour, 1981
34	John Larson	Fe-Zn	Strowd et al., 1981



Map No.	Name	Commodity	Source of Data
35	Snowball	Pb-Zn	Tuchek and Ridenour, 1981
36	Unnamed	Quartz	Tuchek and Ridenour, 1981
37	Little Copper Creek vein	Au-Ag	Strowd et al., 1981
38	Carrier claim	Pb-Ag-Zn	Strowd et al., 1981
39	Copper Bell group	Pb-Ag	Tuchek and Ridenour, 1981
40	Unnamed	Ag	Tuchek and Ridenour, 1981
41	Trudy B No. 1	Pb-Zn	Tuchek and Ridenour, 1981
42	Unnamed	Au-Pb-Cu	Tuchek and Ridenour, 1981
43	Unnamed	Ag-Au-Zn	Tuchek and Ridenour, 1981
44	Unnamed	Limonite staining	Tuchek and Ridenour, 1981
45	Monarch Fraction	Pb-Ag(?) - Zn	Tuchek and Ridenour, 1981
46	John Logan	?	Tuchek and Ridenour, 1981
47	Eagle Bird mine	Ag-Pb-Zn-Cu-Au	Anderson and Wagner, 1946
48	Garfield Antimony (Scorpion)	Ag-Pb-Zn	Strowd et al., 1981
49	Rippeto	Pb-Ag-Zn	Tuchek and Ridenour, 1981
50	Silver Eagle	Ag-Pb-Zn	Tuchek and Ridenour, 1981
51	Smuggler	Ag-Pb-Zn	Tuchek and Ridenour, 1981
52	Carrier	Zn-Ag-Pb-Cu	Strowd et al., 1981
53	Scorpion group	Ag-Pb-Zn	Tuchek and Ridenour, 1981
54	Garfield group	Ag-Pb-Zn	Tuchek and Ridenour, 1981
55	Silver Spar Mining Co.	Ag-Au-Cu-Zn-Pb	Strowd et al., 1981
56	Black Spar	Pb-Ag	Strowd et al., 1981
57	Mutual mine	Pb-Ag	Umpleby, 1917
58	Boundary Gulch prospect	Ag-Pb-Cu-Zn	Tuchek and Ridenour, 1981
59	Grey Eagle	Ag-Pb-Zn	Tuchek and Ridenour, 1981
60	Logan tunnel	Pb-Ag	Tuchek and Ridenour, 1981
61	Champion and Contact	Cu-Pb	Tuchek and Ridenour, 1981
62	Idaho Muldoon mine	Ag-Pb-Zn-Cu	Anderson and Wagner, 1946
63	Solid Muldoon	Pb-Zn-Ag	Anderson and Wagner, 1946



<u>Map No.</u>	<u>Name</u>	<u>Commodity</u>	<u>Source of Data</u>
64	Muldoon mine	Pb-Zn-Ag	Anderson and Wagner, 1946
65	American Mine prospect	Pb-Zn	Anderson and Wagner, 1946
66	Muldoon Ridge	Au	Strowd et al., 1981
67	Boundary and Deep Gulch mines	Ba-Ag-Pb-Zn-Cu	Tuchek and Ridenour, 1981
68	Muldoon Barium	Ba	Tuchek and Ridenour, 1981
69	Unnamed	Pb-Ag	Strowd et al., 1981
70	Lucky Boy	Pb-Ag	Tuchek and Ridenour, 1981
71	Lucky Strike	Ag-Pb-Cu	Nelson and Ross, 1969
72	Garfield mine	Pb-Ag-Zn-Cu	Strowd et al., 1981
73	Silver King mine	Ag-Zn-Pb-W	MILS, 1982; Hustedde et al., 1981
74	Unnamed	Ag-Pb	Tuchek and Ridenour, 1981
75	Porcupine Creek prospect	Pb-Zn	Hustedde et al., 1981
76	Hecla-Carrier	Ag-Pb-Zn	Tuchek and Ridenour, 1981
77	Scorpion Ridge No. 1 and 2	Ag-Pb-Zn	Tuchek and Ridenour, 1981




Legend

CENOZOIC	Qal	Alluvium	Recent	Quaternary
	Qb	Snake River Basalt	Pleistocene	
	Tqm	Quartz monzonite	Eocene	Tertiary
	Tcv	Challis Volcanics		
PALEOZOIC	Mcb	Copper Basin Formation (undifferentiated)		Mississippian

• 10 Location of mine or prospect.

Boundary of Pioneer Mountains GRA.



Mines and Prospects

13. Lead Belt
14. Butte - Antelope
6. Ella
7. Horn Silver
9. Reliance
11. Oxide Lode
10. St. Louis
12. Last Chance
- A. Pandora
- B. Sam and Tam
- C. Antimony
- D. Edna
- E. Silver Tip
- F. Diamond
- G. Wolframite
- H. Hub
- I. Silver Bell
- J. Golden Chariot
- K. Marten
- L. Copper King
- M. Paymaster

Data from: Anderson, (1929).

WGM Inc. Mining and Geological Consultants
Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS

Pioneer Mountains GRA, Idaho
Lava Creek District
Geologic Map

SCALE AS SHOWN
DATE BY: DATE: 3/1983
DRAWN BY: DS1: APRVD: G.F.


FIGURE
11



extends for dozens of feet into the wall rock from the veins, especially in the volcanics. Alteration consists of bleaching accompanied by sericitization and lesser silicification. The alteration zone within several feet of the vein is often impregnated with disseminated pyrite which is occasionally accompanied by galena and sphalerite. The ore occurs in small shoots. Few of the shoots were mineable for more than a couple of hundred feet along strike and most of the deposits were mined for only a few dozen feet along strike (Anderson, 1929). Anderson (1929) considers the mineralization to be Tertiary in age and suggests that it may be related to an unexposed granitic body.

The Copper Basin mining district was active during the 1880s and early 1890s. Total production credited to the district is 92 tons of ore with a recoverable content of 2.33 ounces silver per ton, 5.5% copper, 10.4% lead, and 7.1% zinc (Tuchek and Ridenour, 1981). The most productive mine in the district was the Copper Basin or Reed and Davidson (outside of the Pioneer Mountains GRA) which produced 200,000 pounds of copper. The most productive mine within the GRA apparently was the Star Hope mine (loc. 29, Fig. 10, Table I) which produced about \$50,000 worth of silver-lead ore (Simons, 1981).

Rocks in the vicinity of the Star Hope mine consist of carbonaceous argillite and quartzite with thin interbeds of limestone intruded by rhyolite and granitic dikes. The sedimentary rocks are folded, generally striking northwest and dipping northeast. Quartz veins in these rocks contain galena, tetrahedrite, and chalcopyrite. The sulfides also occur as



fracture fillings within the vein or as breccia fillings when the vein is brecciated (Tuchek and Ridenour, 1981).

The Little Wood River district has been active intermittently since 1881. Production between 1929 and 1964, the period for which records are available, is summarized in Table II. Earlier production from the principal mine in the district, the Solid Muldoon (loc. 63, Fig. 10, Table I), was about \$200,000 worth of silver by 1910. Significant early production amounting to about \$130,000 also came from the Eagle Bird mine (loc. 47). In addition, about 2,400 tons of barite was produced from the Muldoon Barium mine (loc. 68). Total production from the district is estimated to have been worth about \$330,000 (Simons, 1981). The Little Wood River district includes the western quarter of the Pioneer Mountains GRA, and most of the mineral occurrences are concentrated along Muldoon and Garfield Canyons just north of the Friedman Creek WSA.

The mineralization in the Little Wood River mining district occurs in Paleozoic limestones and argillaceous rocks, particularly the Drummond Mine Limestone. The mineralization is often spatially associated with Tertiary quartz monzonite stocks and dikes as can be seen by the concentration of mineral occurrences adjacent to two small areas with surface exposures of quartz monzonite (Fig. 12). The Paleozoic rocks in the district are folded along a northwest-trending regional anticline, and the rocks are extensively fractured. The mineral deposits are in generally irregular but well-defined shoots localized at the intersections of bedding-plane faults and high-angle cross faults. The mineralization consists of galena, sphalerite, silver-bearing minerals, chalcopyrite and locally barite. Quartz is the main






TABLE II
RECORDED METAL PRODUCTION 1929-1964, LITTLEWOOD RIVER DISTRICT

Year	Mine	Orre (t)	Gold (g)	Silver (g)	Copper (kg)	Lead (kg)	Zinc (kg)
1929	United Mines (Garfield group)	9	--	12,177	14	564	---
1938	Eagle Bird	136	--	6,064	95	1,164	---
1943	---do-----	27	--	19,562	45	5,865	---
1944	---do-----	182	31	60,832	294	17,357	5,653
1945	---do-----	336	62	216,158	366	33,897	2,989
1946	---do-----	188	--	44,473	171	11,478	8,402
1947	Eagle Bird	379	62	106,382	503	27,445	15,610
	Unknown	47	--	7,464	93	1,686	1,709
1948	Eagle Bird	647	93	216,571	938	46,580	19,701
1949	---do-----	513	62	158,002	606	41,024	19,958
1950	---do-----	791	93	224,697	1,769	52,527	34,338
1951	Eagle Bird	288	62	52,791	2,525	16,705	14,026
	Rippeto	25	--	9,703	127	4,189	273
	Hecia-Carrier	1	--	902	---	127	---
1952	Eagle Bird	387	218	123,280	400	33,723	13,927
	Prince Pine	1	--	1,187	283	11,66	26
1953	Eagle Bird	149	31	64,408	---	17,172	7,382
1955	Scorpion	1	--	1,306	272	---	---
1956	---do-----	2	31	529	318	---	---
1965	---do-----	8	31	1,710	560	---	---
TOTAL		4,111	776	1,239,117	9,539	311,369	143,994

Data from: Tuckek and Ridenour (1981, p. 259).





Legend

- | | | | |
|--|--|---------------|------------|
| | Alluvium | Recent | Quaternary |
| | Glacial deposits | | |
| | Quartz monzonite | Pleistocene | Tertiary |
| | Dike, sills and small intrusive masses that range from rhyolite to andesite. | | |
| | Challis volcanics | Eocene | Tertiary |
| | Copper Basin Formation | | |
| | Mcbu - Upper Clastic Unit | Mississippian | |
| | Mcbd - Drummond Mine Limestone | | |
| | Mchl - Lower Clastic Unit | | |
| | Mcu - Glide Mountain Plate of Copper Basin (?) Formation. | Devonian | |
| | Milligen Formation | | |




Outline of Friedman Creek WSA



Location of mine, prospect, or occurrence.

For Key to numbers see table 1

Data from: U.S. Geological Survey and U.S. Bureau of Mines, 1981.



WGM Inc. Mining and Geological Consultants
Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS


Pioneer Mountains GRA, Idaho
Little Wood River District
Geologic Map

SCALE: 1"=1 MILE (1:62,500)	FIGURE
DATE BY: DATE: 3/1983	12
DRAWN BY: DSI	APP'D: G.F.



gangue mineral, but minor calcite is also present. The mineralization occurs as massive irregular to tabular-shaped replacement lenses, as disseminations, and as fissure fillings in shear zones or in breccias. Replacement bodies are often localized along bedding planes. The most important mines in the GRA are the Solid Muldoon (loc. 63, Fig. 11), the Eagle Bird (loc. 47), and the Muldoon Barium (loc. 68) properties (Tuchek and Ridenour, 1981). Tuchek and Ridenour (1981) also consider the Idaho Muldoon (loc. 62, Fig. 12), Black Spar (loc. 56, Fig. 12), Drummond (loc. 32, Fig. 10), Muldoon Ridge (loc. 66, Fig. 10), Boundary Gulch (loc. 58, Fig. 12), Champion (loc. 61, Fig. 12), Mutual (loc. 57, Fig. 12), Scorpion (loc. 42, Fig. 12), Garfield (loc. 54, Fig. 12), Silver Eagle (loc. 50, Fig. 12), and Rippeto (loc. 49, Fig. 12) properties to have exploration potential.

At the Solid Muldoon mine (loc. 63, Fig. 12), northwesterly striking quartzites and slates are intruded by porphyry dikes. The sedimentary rocks dip moderately to the northeast. A cross-section of the geology at the Solid Muldoon mine is shown in Figure 13. A dense brecciated sulfide-quartz vein is present immediately above the altered dacite porphyry dike. This zone grades upward into a zone of open-space filling characterized by quartz and minor pyrite. Adjacent to the dike, the argillites are hornfelsed and pyritized (Anderson and Wagner, 1946). The orebodies consisted of replacement deposits localized along bedding planes and generally adjacent to the dikes. Principal minerals include argentiferous galena, pyrite, sphalerite, and chalcopyrite. A few oreshots reportedly contained 70% lead across a thickness of 12 to 20 feet. Tenor of the ore apparently decreased with depth (Tuchek and Ridenour, 1981).




Muldoon Ridge



A. Photograph of Solid Muldoon Mine, looking west. (From Tuck and Ridenour, 1981).

Muldoon Ridge



B. Schematic sketch of geology.

100 50 0 100 FEET

APPROXIMATE SCALE

WGM Inc. Mining and Geological Consultants
Anchorage, Alaska


BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS

Pioneer Mountains GRA, Idaho
Solid Muldoon Mine
Geology

SCALE: AS SHOWN
DATE BY: DATE 3/1983 BY: NHT
DRAWN BY: DBI APPROV: G.F.


FIGURE

13



The Eagle Bird mine (loc. 47, Fig. 12) is in medium- to thick-bedded, dark-blue-gray to black, locally siliceous limestones which strike north-easterly to northwesterly and dip from 30° to 60° northeast. Two principal sets of fractures are present: (1) northwest-trending reverse bedding-plane faults, and (2) westerly to northwesterly high-angle reverse cross faults. The cross faults are marked by locally bleached and limonite-stained zones of fractured rock and gouge. The orebodies occur along or adjacent to two distinct zones of bedding plane faults at or near their intersection with the high-angle cross faults. Galena with lesser amounts of sphalerite, arsenopyrite, complex antimonides, pyrite and local chalcopyrite occurs in massive irregular to tabular shaped replacement bodies. Disseminated mineralization is also present. The lenticular sulfide masses attain thicknesses up to 3 feet and may extend for tens of feet along strike and down-dip. The near-surface orebodies are oxidized. Alteration of the host rock consists mainly of a softening and bleaching of normally resistant limestone. Locally, more intense alteration produces a creamy-white marble (Tuchek and Ridenour, 1981).

The barite at the Muldoon Barium mine (loc. 68, Fig. 12) occurs in three irregular veins ranging from 1.5 to 14 feet thick. One vein may be up to one mile long. A silver-lead-zinc vein occurs nearby (Tuchek and Ridenour, 1981). According to Tuchek and Ridenour (1981), the property was active in 1974 and 4,000 tons of ore were stockpiled. Simons (1981) reports the occurrence of barite talus along with anomalous barium values in geochemical samples from the ridge between Muldoon Canyon and Argosy Creek east of the mine. This area is just west of the boundary of WSA 53-5.




3.3 Known Mineral and Energy Prospects, Occurrences, and Mineralized Areas

A number of other silver-lead-zinc-copper prospects, occurrences, and mineralized areas are known in the Pioneer Mountains GRA. Most of these consist of either vein or replacement mineralization and exhibit the same style of mineralization as the deposits discussed in the previous section; thus, additional discussion of these prospects is unnecessary. These occurrences are tabulated in Table I and shown on Figure 10. A number of the occurrences shown on Figure 10 (locs. 15-20, 30 and 69) were compiled from 15-minute topographic quadrangle maps by Strowd et al. (1981) and little other information is available. Iron Bog Creek which drains three of these occurrences (loc. 18-20) has anomalous lead contents in stream silt samples (Simons, 1981). Three occurrences (locs. 2-4) consist of prospect pits or shafts noted in Bureau of Land Management files (Carroll, pers. comm., 1981), but no other information is known.

Bureau of Land Management internal files (Carroll, pers. comm., 1981) also show two groups of claims (locs. 1 and 5, Fig. 10) which encompass agate occurrences. One of these occurrences, the Espinoza claims (loc. 5), also has some evidence of copper mineralization. Little else is known about these two occurrences.

To evaluate the potential for base-precious metals mineralization along northwest-trending structures which may project into the Friedman Creek WSA, thirty-three stream sediment samples were collected from streams draining the WSA (Appendix III). The analytical data was compared with the data from the Boulder-Pioneer Study Area (Simons, 1981). Using the same threshold



values determined by Simons (1981), eleven samples are anomalous in one or more of the following elements: copper, lead, zinc, and silver (Table III). The anomalies (Fig. 14) occur mainly in valleys with outcropping Paleozoic rocks, although the strong multi-element anomaly at the head of Friedman Creek is underlain by Challis Volcanics. The stream sediment survey outlined eight drainage basins which wholly or partly overlap the WSA and have anomalous lead, zinc, copper, and/or silver contents in their stream sediments. These basins range from slightly less than one-quarter square mile to almost one square mile in area.

There are twelve gravel pits in the eastern part of the Pioneer Mountains GRA (Strowd et al., 1981). The gravel pits are probably utilized on an as-needed basis for local sand and gravel supplies.

3.4 Mining Claims, Leases and Material Sites

A review of BLM claims records current to June 7, 1982 and mineral title plats current to August 12, 1982 shows that there are a total of 1,275 unpatented mining claims and 42 patented mining claims in the Pioneer Mountains GRA (Fig. 15). The majority of the claims are in the Lava Creek and Little Wood River mining districts. About 30 claims are located in or immediately adjacent to the eastern part of the Little Wood River WSA and several claims may overlap the northwest corner of the Friedman Creek WSA. The majority of the patented claims are in the Little Wood River mining district.





TABLE III

ANOMALOUS* GEOCHEMICAL SAMPLES, FRIEDMAN CREEK WSA


Sample No.	Type	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Au (ppm)
28014	Silt	55	45	345	4.0	0.02
28015	Silt	65	50	195	2.2	0.02
28016	Silt	95	50	135	2.2	0.02
28031	Silt	45	45	390	0.2	0.04
28032	Silt	90	60	500	1.2	0.04
28036	Silt	335	335	450	5.0	0.10
28041	Silt	40	175	145	1.0	0.02
28044	Silt	60	55	465	1.0	0.04
28052	Silt	50	170	140	0.8	0.02
28055	Silt	45	35	465	0.4	0.02
28056	Silt	140	60	150	1.0	0.02
28057	Silt	45	40	400	0.4	0.02

* Based on threshold values determined by Simons (1981).



R. 21 E. R. 2 E.

R. 22 E. R. 23 E.



BOUNDARY OF FRIEDMAN WSA (53-5)

THRESHOLD VALUES *

Cu 70 ppm
 Pb 100 ppm
 Zn 300 ppm
 Ag 1.5 ppm
 Au 0.1 ppm

* from: Simons (1981)

T.3N.
 T.2N.

Legend

○ 28050

Stream sediment sample site.



Area of influence of sample
 and anomalous elements.

WGM Inc. Mining and Geological Consultants
 Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT
 REGION 2 NORTHERN ROCKY MOUNTAINS

Pioneer Mountains GRA, Idaho
Geochemical Anomalies
Friedman Creek WSA

SCALE: 1"=1 MILE (1:62,500)


DATA BY: WGM DATE: 3/1983

DRAWN BY: DSI APP'D: G.F.

NOTED

FIGURE


14



EXPLANATION

x Unpatented Mining Claim

▲ Patented Mining Claim



Notes: 1) Number adjacent to symbol indicates number of mining claims in that section.


2) Area of search: Entire GRA

3) Source of Data: BLM Claims Microfiche and MT Plats.

4) Data current to: June 6, 1982.



Approximate Boundary of Wilderness Study Area.



WGM Inc. Mining and Geological Consultants
Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS

Pioneer Mountains, Idaho
Mining Claims Density Map


SCALE: 1"=4 MILES	FIGURE
DATE BY WGM: 1/1983	15
DRAWN BY: DSI	APPROV. BY: G.F.




BLM oil and gas plats current to August 12, 1982 show that much of the eastern half of the Pioneer Mountains GRA is covered by oil and gas leases and lease applications (Fig. 16). Most of the Appendicitis Hill WSA and about 15% of the White Knob Mountains are covered by oil and gas leases (Table IV). About 50% of the White Knob Mountains WSA and about 10% of the Friedman Creek WSA are covered by oil and gas lease applications.

3.5 Mineral and Energy Deposit Types

The most obvious metallic mineral potential in the Pioneer Mountains GRA is for silver-lead-zinc-copper deposits similar to the known occurrences discussed in Section 3.2. Similar deposits are common along the east flank of the Atlanta lobe of the Idaho batholith (Bennett, 1980) and in the fault block ranges east of the batholith (Ruppel, 1978, 1982). Examples are the Bayhorse (McIntyre et al., 1976; Ross, 1937), Gilmore (Ruppel et al., 1979), Lava Creek (Anderson, 1929; Ross, 1931), Little Wood River (Nelson and Ross, 1969; Tuckey and Ridenour, 1981) and the Copper Basin (Tuckey and Ridenour, 1981) districts. The mineralization in these districts consists of irregular veins and replacement bodies in folded and faulted sedimentary and metamorphic rocks and to a lesser extent in intrusive rocks. Quartzite, argillite, limestone and dolomite are common host rock lithologies, however limestone and dolomite are the most abundant hosts (Hobbs, 1968). Argentiferous galena and sphalerite with lesser amounts of pyrite and chalcopyrite are the principal metallic sulfide minerals. The deposits are commonly oxidized at the surface and much of the early production consisted of high-grade secondary silver-lead-zinc ores. The mineralization is associated





WGM Inc. Mining and Geological Consultants
 Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT
 REGION 2 NORTHERN ROCKY MOUNTAINS

Pioneer Mountains, Idaho

Oil and Gas Lease Status

SCALE 1" = 4 MILES		FIGURE 16
DATA BY WGM	DATE 1/1983	
DRAWN BY DSI	APPROV. GF	





TABLE IV
OIL AND GAS LEASES IN WSAs 31-14 AND 31-17


<u>WSA</u>	<u>Lease No.</u>	<u>Owner of Record</u>	<u>Date Issued</u>
31-14	I 16602	Ronan Inc. P.O. Box 1354 Billings, Montana 59103	12-01-81
	I 16697	Ronan Inc. P.O. Box 1354 Billings, Montana 59103	10-01-81
	I 16728	Pioneer Production Corp. P.O. Box 2542 Amarillo, Texas 79189	07-01-81
	I 17034	Wolter Oil Company P.O. Box 8686 Boise, Idaho 83707	10-01-81
	I 17796	Thomas D. Hewitt 915 Gulf Canada Square Calgary, Alberta, Canada	01-01-82
	I 18237	Ronan Inc. P.O. Box 1354 Billings, Montana 59103	05-01-82
31-17	I 18165	Ronan Inc. P.O. Box 1354 Billings, Montana 59103	10-06-81 (applic. filed)
	I 18189	Hewitt Producing Company 915-401 9th Avenue S.W. Calgary, Alberta, Canada T2P 3C5	07-01-82
	I 18190	Hewitt Producing Company 915-401 9th Avenue S.W. Calgary, Alberta, Canada T2P 3C5	10-19-81 (applic. filed)
	I 18194	Hewitt Producing Company 915-401 9th Avenue S.W. Calgary, Alberta, Canada T2P 3C5	10-19-81 (applic. filed)
53-5	I 18440	John R. Warne 3/8 interest Irvin Kronzler " Robert C. Balson " Box 1324 Billings, Montana 59103	12-24-81



with Tertiary intrusive bodies (Bennett, 1980). These rocks are often hydrothermally altered and contain disseminated sulfides (Ruppel, 1982).

Tungsten occurs in a few of the silver-lead-zinc districts mentioned above as an accessory. Cook (1955) delineates a northwest-trending belt of contact tungsten deposits which runs from the Lava Creek mining district northwest to the Big Creek mining district in northern Valley County. The tungsten occurs as scheelite or powellite in calcareous rocks which have been thermally metamorphosed by nearby intrusive bodies.

Most authors who have discussed the genesis of the metallic mineral deposits typical of south-central Idaho emphasize the association of the mineral deposits with intrusive rocks and structures. Ruppel (1978, 1982) concludes that most of the major mineral deposits in east-central Idaho were genetically associated with neck-like, central stocks which rose along steep reverse faults. These stocks fed magma laterally along the imbricate thrust faults resulting in the formation of sheet-like bodies (Ruppel, 1982). Bennett (1980) proposes that convective hydrothermal systems were formed around epizonal Tertiary plutons. Such a concept is supported by Hall et al. (1978) in their geochemical study of the Wood River district. They conclude that the mineral deposits in the district formed by leaching of metals from Paleozoic country rocks and redeposition in favorable structural settings. The distribution of most of the known mineralization in the Pioneer Mountains GRA does appear to be spatially associated with Tertiary intrusives; thus, similar undiscovered deposits associated with buried intrusives may exist at depth. Aeromagnetic data covering the northwest



part of the GRA, including WSA 53-4 and 53-5, shows a strong northwest-trending magnetic high which includes the Muldoon stock (Mabey, 1981). The gentle slope of the southeast end of the anomaly suggests that the nose of the stock may extend beneath the northwest part of WSA 53-5. The geochemical anomalies in the area between Muldoon Canyon and Friedman Creek are within the area that may be underlain by the nose of the pluton. A second magnetic peak occurs on the northwest-trending magnetic high near the confluence of Muldoon and Copper Creeks. Mabey (1981) interprets this peak to be a shallow stock. Both stocks are interpreted to be parts of a much larger northwest-trending pluton underlying the large magnetic high (Mabey, 1981).

The sedimentary rocks underlying the Pioneer Mountains GRA were deposited in a continental margin and continental rise and continental rise setting. Major metallic mineral deposit types expected in this setting are carbonate or sandstone-hosted lead-zinc deposits and metalliferous black shale/argillite deposits (Mitchell and Garson, 1981).

The carbonate or sandstone-hosted lead-zinc deposits in this setting are collectively known as Mississippi Valley-type mineralization which is generally characterized by the following:

1. simple sulfide mineralogy
2. generally hosted by dolomites, less commonly by sandstones or limestones
3. absence of nearby intrusive rocks
4. mineralization related to local facies changes in host rock

1. The first part of the report is a summary of the work done during the year.

2. The second part is a detailed account of the work done during the year.

3. The third part is a summary of the work done during the year.

4. The fourth part is a summary of the work done during the year.

5. The fifth part is a summary of the work done during the year.

6. The sixth part is a summary of the work done during the year.

7. The seventh part is a summary of the work done during the year.

8. The eighth part is a summary of the work done during the year.

9. The ninth part is a summary of the work done during the year.

10. The tenth part is a summary of the work done during the year.

11. The eleventh part is a summary of the work done during the year.

12. The twelfth part is a summary of the work done during the year.

13. The thirteenth part is a summary of the work done during the year.

14. The fourteenth part is a summary of the work done during the year.

15. The fifteenth part is a summary of the work done during the year.

16. The sixteenth part is a summary of the work done during the year.

17. The seventeenth part is a summary of the work done during the year.

18. The eighteenth part is a summary of the work done during the year.

19. The nineteenth part is a summary of the work done during the year.

20. The twentieth part is a summary of the work done during the year.

21. The twenty-first part is a summary of the work done during the year.

22. The twenty-second part is a summary of the work done during the year.

23. The twenty-third part is a summary of the work done during the year.

24. The twenty-fourth part is a summary of the work done during the year.

25. The twenty-fifth part is a summary of the work done during the year.

26. The twenty-sixth part is a summary of the work done during the year.

27. The twenty-seventh part is a summary of the work done during the year.

28. The twenty-eighth part is a summary of the work done during the year.

29. The twenty-ninth part is a summary of the work done during the year.


30. The thirtieth part is a summary of the work done during the year.

5. districts related to basal margins
6. relatively simple structure at time of deposition

The Paleozoic ages of leads from the Wood River district (Hall et al., 1978), the general characteristics of some of the lead-zinc mineralization in east-central Idaho, and the presence of stratabound lead and copper mineralization in Cambrian rocks in southeastern Idaho (Mansfield, 1927) suggest regional potential for Mississippi Valley-type mineralization in the older continental margin setting of east-central Idaho.

Although the allochthonous nature of the terrane within the Pioneer Mountains GRA obscures the facies relationships among the sediments, both breccias and reef complexes, which are excellent ore hosts elsewhere, may be present in the Paleozoic carbonates. If Mississippi Valley-type lead-zinc mineralization is present in east-central Idaho, then some of the known lead-zinc occurrences may represent remobilization of undiscovered mineralization during later geologic processes, particularly via ground water migration along fractures and faults and via hydrothermal convection due to igneous intrusion.

Silver-lead-zinc mineralization is sometimes associated with argillitic turbidite sequences. An example of this type of mineralization occurs in the eastern Selwyn basin in the Yukon Territory where stratiform silver-lead-zinc mineralization occurs in a sequence of argillites, limestones, sandstones and conglomerates. Barite is often present as an accessory mineral and barium halos commonly surround the centers of lead-zinc-silver mineralization (Carne, 1976). Simons (1981) did not find the black shales/



argillites in the Boulder-Pioneer study area to be notably enriched in metals, but he did find that the lower clastic unit of the Copper Basin Formation contains unusually large amounts of barium. This suggests that some potential for stratiform silver-lead-zinc mineralization may exist in the Late Devonian and Early Mississippian argillite sequences.

There are no known uranium occurrences in the Pioneer Mountains GRA. A hydrogeochemical and stream geochemical sampling program was conducted in the Hailey NTMS Quadrangle, but only the analytical results from this program have been released. Without location maps, the data could not be used in this study. However in the Beaverhead Range, the McGowan Creek Formation was found to be uraniferous (Wodzicki and Krason, 1981). This unit is present in the GRA, but no data is available on the uranium content of the McGowan Creek Formation within the Pioneer and White Knob Mountains.

The Pioneer Mountains GRA is a part of the Cordilleran fold-thrust belt. Major hydrocarbon reserves are present 175 miles southeast of the GRA in western Wyoming where three giant oil fields (Lamb, 1980; McCaslin, 1981) and 12 other fields are currently being developed. Some of the latter may be classed in the giant ranks when fully developed. Production within this area is from several horizons including Ordovician, Devonian, Mississippian, Pennsylvanian, Triassic, and Jurassic units (McCaslin, 1980, 1981). In the Pioneer Mountains GRA most of these Paleozoic units have equivalents, but Triassic and Jurassic rocks are not present. Production is also known from the Thrust belt in west-central Montana, approximately 365 miles north of the Pioneer Mountains GRA. Several geologists have indicated that the

The first of these is the fact that the
the second is the fact that the
the third is the fact that the

the fourth is the fact that the
the fifth is the fact that the
the sixth is the fact that the


the seventh is the fact that the
the eighth is the fact that the
the ninth is the fact that the

the tenth is the fact that the
the eleventh is the fact that the
the twelfth is the fact that the

the thirteenth is the fact that the
the fourteenth is the fact that the
the fifteenth is the fact that the

Thrust belt structural province extending from southeastern Idaho and southwestern Wyoming continues into east-central Idaho and southwestern Montana (King, 1969; Blackstone, 1977; Ruppel et al., 1981). Potential hydrocarbon source beds in the Pioneer Mountains GRA are the dark shales of the Milligen and McGowan Creek Formations, argillaceous limestones and calcareous shales of the Jefferson Formation, and dark clayey-silty limestones of the South Creek Formation. All of these units except the South Creek Formation have had hydrocarbon analyses made on them from localities in east-central Idaho or southwestern Montana. A petroleum company had samples of the McGowan Creek Formation from the White Knob and Lost River Ranges analyzed. The results of these analyses indicate that the unit has a mature, very poor oil, good to excellent wet-gas condensate source character (Nance Petroleum, pers. comm., 1982). Minor source beds could occur in the shaly limestones of the White Knob and Wood River Formations; however, insufficient studies of these units have been made to verify the hydrocarbon source potential.

Potential reservoir beds in the Paleozoic section include sucrosic dolomite in the Ordovician should it be present at depth; vuggy Silurian Laketown Dolomite, fractured carbonates of Ordovician and younger age, and sandstones of late Paleozoic age. Samples of vuggy Laketown Dolomite collected in the northern part of the White Knob Mountains would make excellent hydrocarbon reservoirs given the proper structural conditions (Nance Petroleum, pers. comm., 1982). Insufficient studies have been made of the units underlying the Pioneer Mountains GRA to quantify reservoir characteristics of any of the formations underlying the area.



Structural traps in the Pioneer Mountains GRA are not defined at present. The presence imbricated thrust plates in the area (Skipp and Hait, 1977) as well as the existence of anticlinal folds, (Nelson and Ross, 1969) and high angle faulting (Ruppel, 1982) suggests that excellent structural traps should be present in the subsurface of the Pioneer Mountains GRA. In addition, magnetic and seismic data suggests that structural traps are present in WSA 31-14 and 31-17 (Nance Petroleum, pers. comm., 1982).

Considering that commercial production exists in the Thrust belt both sides of the Pioneer Mountains GRA and that the GRA is part of this major structural belt, oil and gas potential in the GRA is good. However, until the area is tested by drilling it will remain unknown whether or not commercial hydrocarbon deposits are present.

There are no known hot springs or other geothermal manifestations in the Pioneer Mountains GRA; therefore, the geothermal potential of the area must be evaluated by analogy based on the regional setting. The Pioneer Mountains GRA is within the Central Idaho Basin and Range geothermal province which occupies the area between the Idaho batholith on the west, the Montana Basin and Range province on the east, and the Snake River Plains on the south. The area is distinguished from the Montana Basin and Range in that the bedrock of the province is predominately Precambrian and Paleozoic sedimentary rocks which have been extensively thrust faulted. Except for a few outliers of basalt along the margin of the Snake River Plain, the Eocene Challis Volcanics are the youngest volcanic rocks exposed in this province. Normal faulting appears to have been active until very recent times. In addition, a seismic trend extending westward from Yellowstone National Park

passes through the center of the province. Another zone of seismic activity is present just west of the Central Idaho Basin and Range geothermal province and an east-west trend of seismic activity is also present along the Centennial Mountains on the Montana border just east of the province (Smith, 1978). The hot spring frequency in the Central Idaho Basin and Range geothermal province is somewhat lower than in the Montana and Basin Range province; however, the Central Idaho basin and Range province is virtually unknown on even a reconnaissance geothermal or geophysical basis. Typical heat flow values are probably the same as the Montana Basin and Range province, but very little data is available. Due to the abundance of carbonate rocks in the province, the hydrology is probably more complicated than in the Montana Basin and Range province where less permeable granitic and metamorphic rocks predominate. However, it is not possible to predict the influence that bedrock hydrology might have on the location of geothermal systems in the Central Idaho Basin and Range province using existing data.

There are no wells with thermal gradient information in the immediate vicinity of the Pioneer Mountains GRA. South of Timbered Dome, a 360 foot deep well has a surface temperature of 43.5°C. Wells at Butte City, 10 miles southeast of Appendicitis Hill, may have temperatures as high as 76°C at a depth of 86 feet. These wells are along the boundary between the Snake River Plain and the Central Idaho Basin and Range geothermal provinces (Mitchell et al., 1980). The geothermal setting northwest of this boundary is probably typical of that throughout the Central Idaho Basin and Range geothermal province.


3.6 Mineral and Energy Economics

There are good gravel roads throughout most of the Pioneer Mountains GRA and adequate water sources. These factors combine to make mineral development in the GRA economically attractive given a suitable deposit.


The rise in the price of silver in the late 1970s spurred considerable exploration interest. U.S. market demand has steadily increased as more industrial uses for silver are found (Rosta, 1982). Silver deposits formerly considered to be low grade are now economically attractive at present prices. Additionally, recoverable silver values in base metal ores, e.g. lead or copper, now make a substantial contribution to mine profits.

With the increase in the price imported oil in the past decade, the deregulation of domestic natural gas prices, and increasing government emphasis on energy self sufficiency, exploration and development of domestic oil and gas resources has proceeded at an accelerated pace. Natural gas reserves discovered in the late 1950s and 1960s were often not developed because of a lack of market (Heany, 1961). However, under present conditions drilling is taking place with a goal of near-term production.

The economic feasibility of geothermal resource development is determined by its distance to market and its temperature. Long-distance transportation of lower temperature geothermal energy is not feasible whereas for electrical grade resources long transportation distances are feasible. Based on present requirements for the use of hot fluids in electrical generating techniques, geothermal systems with temperatures of less than 150°C do not



have significant potential for electrical exploitation. However, geothermal resources with temperatures less than 150°C do have significant potential for low and intermediate temperature geothermal utilization for space heating, material processing, etc. if their minimum temperature exceeds 40°C. At the lower end of the spectrum, as the energy content of the resource becomes less, or the drilling depth necessary for exploitation becomes greater, there is a very ill-defined cutoff. For example, shallow ground water temperatures on the order of 10-20°C can be used for heat pump applications, and in some cases these are considered geothermal resources. However, in this evaluation, a lower temperature than approximately 40°C is considered uneconomic as a geothermal resource.



4.0 LAND CLASSIFICATION FOR GEM RESOURCES POTENTIAL

4.1 Explanation of Classification Scheme

In the following sections the land in the WSAs within the Pioneer Mountains GRA is classified for geology, energy and mineral (GEM) resources potential. The classification scheme used is shown in Table V. Use of this scheme is specified in the contract under which WGM prepared this report.

The evaluation of resource potential and integration into the BLM classification system has been done using a combination of simple subjective and complex subjective approaches (Singer and Mosier, 1981) to regional resources assessment. The simple subjective approach involves the evaluation of resources based on the experience and knowledge of the individuals conducting the evaluations. The complex subjective method involves use of rules, i.e. geologic inference, based on expert opinion concerning the nature and importance geologic relationships associated with mineral and energy deposits (Singer and Mosier, 1981).

The GEM evaluation is the culmination of a series of tasks. The nature and order of the tasks was specified by the BLM, however they constitute the general approach by which most resource evaluations of this type are conducted. The sequence of work was: (1) data collection, (2) compilation, (3) evaluation, and (4) report preparation. Two days of field work were done in the Pioneer Mountains GRA.




TABLE VI


BUREAU OF LAND MANAGEMENT GEM RESOURCES LAND CLASSIFICATION SYSTEM

CLASSIFICATION SCHEME

1. The geologic environment and the inferred geologic processes do not indicate favorability for accumulation of mineral resources.
2. The geologic environment and the inferred geologic processes indicate low favorability for accumulation of mineral resources.
3. The geologic environment, the inferred geologic processes, and the reported mineral occurrences indicate moderate favorability for accumulation of mineral resources.
4. The geologic environment, the inferred geologic processes, the reported mineral occurrences, and the known mines or deposits indicate high favorability for accumulation of mineral resources.

LEVELS OF CONFIDENCE

- A. The available data are either insufficient and/or cannot be considered as direct evidence to support or refute the possible existence of mineral resources within the respective area.
- B. The available data provide indirect evidence to support or refute the possible existence of mineral resources.
- C. The available data provide direct evidence, but are quantitatively minimal to support or refute the possible existence of mineral resources.
- D. The available data provide abundant direct and indirect evidence to support or refute the possible existence of mineral resources.




Each WSA is classified for locatable, leasable, and saleable resources potential.

Locatable minerals are those which are locatable under the General Mining Law of 1872, as amended, and the Placer Act of 1870, as amended. Minerals which are locatable under these acts include metals, ores of metals, non-metallic minerals such as asbestos, barite, zeolites, graphite, uncommon varieties of sand, gravel, building stone, limestone, dolomite, pumice, pumicite, clay magnesite, silica sand, etc. (Maley, 1983).

Leasable resources include those which may be acquired under the Mineral Leasing Act of 1920 as amended by the Acts of 1927, 1953, 1970, and 1976. Materials covered under this Act include: asphalt, bitumen, borates of sodium and potassium, carbonates of sodium and potassium, coal, natural gas, nitrates of sodium and potassium, oil, oil shale, phosphate, silicates of sodium and potassium, sulfates of sodium and potassium, geothermal resources, etc. (Maley, 1983).

Saleable resources include those which may be acquired under the Materials Act of 1947 as amended by the Acts of 1955 and 1962. Included under this Act are common varieties of sand, gravel, stone, cinders, pumice, pumicite, clay, limestone, dolomite, peat and petrified wood (Maley, 1983).



4.2 Classification of the Appendicitus Hill (31-14) WSA

4.2.1 Locatable Resources


4.2.1a Metallic Minerals. The Appendicitus Hill WSA (1a, Fig. 17) is classified as having low to moderate favorability for metallic mineral resources, but the available data is insufficient for an adequate evaluation (2A-3A). The low confidence rating is assigned because of the low level of detail of published geologic mapping and the lack of geochemical and geophysical data.

4.2.1b Uranium and Thorium. WSA 31-14 (1b, Fig. 17) is classified as having low favorability for uranium and thorium based on inadequate data (2A). No occurrences are known, but the data base is sparse. The classification should be higher if the McGowan Creek Formation is found to be uraniferous.

4.2.1c Non-Metallic Minerals. All of WSA 31-14 (1c, Fig. 17) is classified as having low to moderate potential for barite, but the available data is insufficient for an adequate evaluation (2A-3A). The low level of confidence is assigned because of the low level of detail of available geologic mapping and the lack of geochemical and geophysical data.

4.2.2 Leasable Resources

4.2.2a Oil and Gas. All of WSA 31-14 (1 and 2, Fig. 18) is classified as having moderate favorability for oil and gas based on limited direct



Pioneer Mountains GRA, Idaho
Legend For Figure 17

BLM LAND CLASSIFICATION SYSTEM
FOR GEM RESOURCES


CLASSIFICATION SCHEME

1. THE GEOLOGIC ENVIRONMENT AND THE INFERRED GEOLOGIC PROCESSES DO NOT INDICATE FAVORABILITY FOR ACCUMULATION OF MINERAL RESOURCES.
2. THE GEOLOGIC ENVIRONMENT AND THE INFERRED GEOLOGIC PROCESSES INDICATE LOW FAVORABILITY FOR ACCUMULATION OF MINERAL RESOURCES.
3. THE GEOLOGIC ENVIRONMENT, THE INFERRED GEOLOGIC PROCESSES, AND THE REPORTED MINERAL OCCURRENCES INDICATE MODERATE FAVORABILITY FOR ACCUMULATION OF MINERAL RESOURCES.
4. THE GEOLOGIC ENVIRONMENT, THE INFERRED GEOLOGIC PROCESSES, THE REPORTED MINERAL OCCURRENCES, AND THE KNOWN MINES OR DEPOSITS INDICATE HIGH FAVORABILITY FOR ACCUMULATION OF MINERAL RESOURCES.


LEVELS OF CONFIDENCE


- A. THE AVAILABLE DATA ARE EITHER INSUFFICIENT AND/OR CANNOT BE CONSIDERED AS DIRECT EVIDENCE TO SUPPORT OR REFUTE THE POSSIBLE EXISTENCE OF MINERAL RESOURCES WITHIN THE RESPECTIVE AREA.
- B. THE AVAILABLE DATA PROVIDE INDIRECT EVIDENCE TO SUPPORT OR REFUTE THE POSSIBLE EXISTENCE OF MINERAL RESOURCES.
- C. THE AVAILABLE DATA PROVIDE DIRECT EVIDENCE, BUT ARE QUANTITATIVELY MINIMAL TO SUPPORT OR REFUTE THE POSSIBLE EXISTENCE OF MINERAL RESOURCES.
- D. THE AVAILABLE DATA PROVIDE ABUNDANT DIRECT AND INDIRECT EVIDENCE TO SUPPORT OR REFUTE THE POSSIBLE EXISTENCE OF MINERAL RESOURCES.

EXPLANATION



- a) Metallic Minerals
b) Uranium and Thorium
c) Non-Metallic Minerals





Approximate Boundary of Wilderness Study Area.

WGM Inc. Mining and Geological Consultants
Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS

Pioneer Mountains, Idaho
Wilderness Study Area
Land Classification
Locatable Resources


SCALE: 1" = 4 MILES (1:250,000)

DATA BY: WGM DATE: 9/1982

DRAWN BY: DSI APPROVED BY: G.F.


FIGURE

17



- a) oil and gas
- b) Geothermal: high temperature (H), Low temperature (L)
- c) Sodium and Potassium
- d) others: Asphalt (As) bitumen (bt), phosphate (ph), No specific commodity designation indicates that the rating applies to all of the above.






Approximate Boundary of Wilderness Study Area.

WGM Inc. Mining and Geological Consultants
Anchorage, Alaska

BLM GEM RESOURCES ASSESSMENT
REGION 2 NORTHERN ROCKY MOUNTAINS

Pioneer Mountains, Idaho
Wilderness Study Area
Land Classification
Leasable Resources

SCALE: 1" = 4 MILES (1:250,000)	FIGURE
DATA BY: WGM DATE: 9/1982	18
DRAWN BY: DSI	
APP'D: G.F.	




evidence (3C). The basis of this classification is the structural setting of the WSA including excellent potential for the development of traps, indications of subsurface structures from private geophysical data, the presence of potential hydrocarbon source and reservoir beds in the stratigraphic section, and a favorable thermal history of the source rocks.

4.2.2b Geothermal. Most of WSA 31-14 (1, Fig. 18) is classified as being unfavorable for geothermal resources based on indirect evidence (1B). The portions of WSA 31-14 along Antelope Creek and Lost River Valleys (2, Fig. 18) are classified as having low favorability for geothermal resources based on indirect evidence (2B). The 1B classification is based on analogy with similar areas in the Idaho Basin and Range geothermal province and the high elevations present combined with the absence of major faults or lineaments. The 2B classification is based on proximity to the northern margin of the Snake River Plain, the existence of a range-bounding fault along the east side of the WSA, and the presence of a major lineament along Antelope Creek.

4.2.3c Sodium and Potassium. All of WSA 31-14 (1 and 2, Fig. 18) is classified as having low favorability for sodium and potassium based on indirect evidence (2B). The basis of the classification is the generally unfavorable geologic environment present in the WSA and the lack of known occurrences in the region.

4.2.3d Other. WSA 31-14 (1 and 2, Fig. 18) is classified as having low favorability for other leasable resources, including phosphate, bitumen, and asphalt based on indirect evidence (2B). The basis of this classification



is the lack of known occurrences in the area and the generally unfavorable geologic environment of the WSA.

4.2.3 Saleable Resources


WSA 31-14 (1, Fig. 19) is classified as having moderate favorability for common varieties of limestone based on indirect evidence (3B). The basis of the classification is the dominance of carbonate rocks in the stratigraphic section. The lack of detailed geologic mapping in the area precludes a more site-specific classification.

4.3 Classification of the White Knob Mountains (31-17) WSA


4.3.1 Locatable Resources

4.3.2a Metallic Minerals. All of WSA 31-17 (2a, Fig. 17) is classified as having low to moderate favorability for metallic mineral resources, but the available data is insufficient for an adequate evaluation (2A-3A). The low confidence rating is assigned because of the low level of detail of published geologic mapping and the lack of geochemical and geophysical data.

4.3.1b Uranium and Thorium. WSA 31-17 (2b, Fig. 17) is classified as having low favorability for uranium and thorium based on inadequate evidence (2A). No occurrences are known, but the classification should be higher if uraniferous McGowan Creek Formation is present in the WSA.







4.3.1c Non-metallic Minerals. WSA 31-17 (2c, Fig. 17) is classified as having low to moderate potential for barite, but the available data are insufficient for an adequate evaluation (2A-3A). The low level of confidence is assigned because of the low level of detail of available geologic mapping and the lack of geochemical and geophysical data.

4.3.2 Leasable Resources

4.3.2a Oil and Gas. All of WSA 31-17 (3, Fig. 18) is classified as having moderate favorability for oil and gas based on limited direct evidence (3C). The basis of this classification is the structural setting of the WSA including excellent potential for development of traps, indications of subsurface structures in private geophysical data, the presence of hydrocarbon source and reservoir beds in the stratigraphic section, and the favorable thermal history of the source rocks.

4.3.2b Geothermal. All of the area of WSA 31-17 (3, Fig. 18) is classified as being unfavorable for geothermal resources based on indirect evidence (1B). The classification is based analogy with similar areas with the Idaho Basin and Range province. Negative factors include generally high elevations and an absence of major faults or lineaments.

4.3.2c Sodium and Potassium. All of WSA 31-17 (3, Fig. 18) is classified as having low favorability for sodium and potassium based on indirect evidence (2B). The basis of the classification is the generally unfavorable geologic environment of the WSA and the lack of known occurrences in the region.

The first part of the report deals with the general situation of the country. It is a very interesting and informative study of the country's development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country.

The second part of the report deals with the economic situation of the country. It is a very interesting and informative study of the country's economic development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country.

The third part of the report deals with the social situation of the country. It is a very interesting and informative study of the country's social development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country.

The fourth part of the report deals with the political situation of the country. It is a very interesting and informative study of the country's political development. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country.

The fifth part of the report deals with the future of the country. It is a very interesting and informative study of the country's future. The author has done a great deal of research and has gathered a wealth of material. The report is well written and is a valuable contribution to the study of the country.

4.3.2d Other. WSA 31-17 (3, Fig. 18) is classified as having low favorability for other leasable resources, including phosphate, bitumen, and asphalt based on indirect evidence (2B). The basis of this classification is the lack of known occurrences in the area and the generally unfavorable geologic environment of the WSA.

4.3.3 Saleable Resources

WSA 31-17 (2, Fig. 19) is classified as having moderate favorability for common varieties of limestone based on indirect evidence (3B). The basis of the classification is the dominance of carbonate rocks in the stratigraphic section. The lack of detailed geologic mapping in the area precludes a more site-specific classification.

4.4 Classification of the Little Wood River (53-4) WSA

4.4.1 Locatable Resources

4.4.1a Metallic Minerals. WSA 53-4 (3, Fig. 17) is classified as having moderate favorability for metallic minerals based on limited direct evidence (3C). The basis of this classification is the favorable geologic setting of the WSA, the presence of known mineral occurrences, and favorable geophysical data.

4.4.1b Uranium and Thorium. WSA 53-4 (3b, Fig. 17) is classified as having low favorability for uranium and thorium based on indirect evidence (2B).

The basis for the classification is the absence of known occurrences.

the first of these is the fact that the
the second is the fact that the
the third is the fact that the

the fourth is the fact that the
the fifth is the fact that the
the sixth is the fact that the

the seventh is the fact that the
the eighth is the fact that the

the ninth is the fact that the
the tenth is the fact that the
the eleventh is the fact that the
the twelfth is the fact that the

the thirteenth is the fact that the
the fourteenth is the fact that the
the fifteenth is the fact that the
the sixteenth is the fact that the

4.4.1c Non-Metallic Minerals. All of WSA 53-4 (3c, Fig. 17) is classified as having moderate potential for barite based on indirect evidence (3B). The low level of confidence is assigned because of the low level of detail of the available geological mapping and the lack of geochemical data.

4.4.2 Leasable Resources

4.4.2a Oil and Gas. WSA 53-4 (4, Fig. 18) is classified as having moderate favorability for oil and gas based on limited direct evidence (3C). The basis of this classification is the favorable structural setting of the WSA including excellent potential for development of traps, indications of subsurface structures in private geophysical data; the presence of hydrocarbon source and reservoir beds in the stratigraphic section, and the favorable thermal history of the source rocks.

4.4.2b Geothermal. WSA 53-4 (4b, Fig. 18) is classified as being unfavorable for geothermal resources based on indirect evidence (1B). The basis of the classification is by analogy with similar areas in the Idaho Basin and Range geothermal province. The WSA is mainly at high elevations and major faults or lineaments are absent.

4.4.2c Sodium and Potassium. WSA 53-4 (4c, Fig. 18) is classified as having low favorability for sodium and potassium based on indirect evidence (2B). The basis of the classification is the generally unfavorable geologic environment of the WSA and the lack of known occurrences in the region.

1. The first part of the document is a letter from the President of the United States to the Congress, dated January 1, 1861.

2. The second part is a report from the Secretary of the Treasury, dated January 1, 1861.

3. The third part is a report from the Secretary of the Interior, dated January 1, 1861.

4. The fourth part is a report from the Secretary of the Navy, dated January 1, 1861.

5. The fifth part is a report from the Secretary of the War, dated January 1, 1861.

6. The sixth part is a report from the Secretary of the State, dated January 1, 1861.

7. The seventh part is a report from the Secretary of the Agriculture, dated January 1, 1861.

8. The eighth part is a report from the Secretary of the Commerce, dated January 1, 1861.

9. The ninth part is a report from the Secretary of the Education, dated January 1, 1861.

10. The tenth part is a report from the Secretary of the Public Works, dated January 1, 1861.

11. The eleventh part is a report from the Secretary of the Public Lands, dated January 1, 1861.

12. The twelfth part is a report from the Secretary of the Public Debt, dated January 1, 1861.

13. The thirteenth part is a report from the Secretary of the Public Buildings, dated January 1, 1861.

14. The fourteenth part is a report from the Secretary of the Public Printing, dated January 1, 1861.

15. The fifteenth part is a report from the Secretary of the Public Works, dated January 1, 1861.

16. The sixteenth part is a report from the Secretary of the Public Lands, dated January 1, 1861.

17. The seventeenth part is a report from the Secretary of the Public Debt, dated January 1, 1861.

18. The eighteenth part is a report from the Secretary of the Public Buildings, dated January 1, 1861.

19. The nineteenth part is a report from the Secretary of the Public Printing, dated January 1, 1861.

20. The twentieth part is a report from the Secretary of the Public Works, dated January 1, 1861.

21. The twenty-first part is a report from the Secretary of the Public Lands, dated January 1, 1861.

22. The twenty-second part is a report from the Secretary of the Public Debt, dated January 1, 1861.

23. The twenty-third part is a report from the Secretary of the Public Buildings, dated January 1, 1861.

24. The twenty-fourth part is a report from the Secretary of the Public Printing, dated January 1, 1861.

25. The twenty-fifth part is a report from the Secretary of the Public Works, dated January 1, 1861.

26. The twenty-sixth part is a report from the Secretary of the Public Lands, dated January 1, 1861.

27. The twenty-seventh part is a report from the Secretary of the Public Debt, dated January 1, 1861.

28. The twenty-eighth part is a report from the Secretary of the Public Buildings, dated January 1, 1861.

29. The twenty-ninth part is a report from the Secretary of the Public Printing, dated January 1, 1861.

30. The thirtieth part is a report from the Secretary of the Public Works, dated January 1, 1861.

4.4.2d Other. WSA 53-4 (4d, Fig. 18) is classified as having low favorability for other leasable resources, including phosphate, bitumen, and asphalt based on indirect evidence (2B). The basis of this classification is the lack of known occurrences in the area and the generally unfavorable geologic environment of the WSA.

4.4.3 Saleable Resources


The entire area of WSA 53-4 (3, Fig. 19) is classified as having low favorability for saleable resources based on indirect evidence (2B).

4.5 Classification of the Friedman Creek (53-5) WSA

4.5.1 Locatable Resources

4.5.1a Metallic Minerals. WSA 53-5 (4a, Fig. 17) is classified as having high favorability for metallic minerals based on abundant direct evidence (4D). The basis of this classification is the immediate proximity and geologic similarity of the WSA to an adjacent mineral district, the trend of mineralized structures into the WSA, and the numerous geochemical anomalies present in the WSA.

4.5.1b Uranium and Thorium. WSA 53-5 (4b, Fig. 17) is classified as having low favorability for uranium and thorium resources based on indirect evidence (2B). The basis for this classification is the absence of known occurrences.




4.5.1c Non-Metallic Minerals. WSA 53-5 (4c, Fig. 17) is classified as having moderate favorability for barite based on limited direct evidence (3C). The basis of this classification is the presence of barite mineralization and barium geochemical anomalies adjacent to the WSA, and the similarity of the geology within the WSA to that of the area surrounding the Muldoon barium deposits.

4.5.2 Leasable Resources

4.5.2a Oil and Gas. WSA 53-5 (4, Fig. 18) is classified as having moderate favorability for oil and gas based on limited direct evidence (3C). The basis of this classification is the structural setting of the WSA including excellent potential for development of traps, indications of subsurface structures in private geophysical data, the presence of hydrocarbon source and reservoir beds in the stratigraphic section, and the favorable thermal history of the source rocks.

4.5.2b Geothermal. All of WSA 53-5 (5, Fig. 18) is classified as being unfavorable for geothermal resources based on indirect evidence (1B). The basis of the classification is by analogy with similar areas in the Idaho Basin and Range province. The area is mostly at high elevations and major faults or lineaments are absent.


4.5.2c Sodium and Potassium. WSA 53-5 (5, Fig. 18) is classified as having low favorability for sodium and potassium based on indirect evidence (2B). The basis of the classification is the generally unfavorable geologic environment of the WSA and the lack of known occurrences in the region.



4.5.3d Other. WSA 53-5 (5, Fig. 18) is classified as having low favorability for other leasable resources, including phosphate, bitumen, and asphalt based on indirect evidence (2B). The basis of this classification is the lack of known occurrences in the area and the generally unfavorable geologic environment of the WSA.

4.5.4 Saleable Resources

The entire area of WSA 53-5 (4, Fig. 19) is classified as having low favorability for saleable resources based on indirect evidence (2B).



5.0 RECOMMENDATIONS FOR FURTHER WORK


The eastern part of the Pioneer Mountains GRA needs to be mapped at a minimum scale of 1:62,500. Available mapping is at reconnaissance scales and was completed at a time when the stratigraphy was not as well understood as today. More detailed mapping of the area would improve the evaluation for oil and gas, metallic minerals and geothermal resources.

The NURE data from the Idaho Falls Quadrangle should be integrated into the data base upon release of the report.

Geochemical stream sediment samples should be collected from the streams draining WSAs 31-14, 31-17 and 53-4 to increase the data base and improve the level of confidence in the mineral assessments. Samples should be collected at a density of 1 to 2 samples per square mile. The samples should be analyzed for copper, lead, zinc, molybdenum, and uranium. During this program the McGowan Creek Formation should also be sampled for uranium.


The base data available on WSA 53-5 are sufficient. More specific evaluation should be completed. This work should include analyses of the samples collected for barium and molybdenum and geological mapping of the WSA and the adjacent Muldoon Canyon area at a scale of 1:24,000. The mapping should be accompanied by detailed rock geochemical sampling.

The hydrocarbon assessment would be upgraded by the completion of the following:




1. Potential hydrocarbon source beds within the Pioneer Mountains GRA should be sampled and analyzed for hydrocarbons. The results would indicate the types of hydrocarbons that might be expected within the area. It would also provide a thermal maturity history of the area.
2. Analyses of potential reservoir beds should be made. These studies will aid in the location of more favorable traps.
3. Conodont color alteration index studies should be made of all Paleozoic carbonates in the Pioneer Mountains GRA. This will provide a thermal maturation history of the Paleozoic stratigraphic section and help indicate the types of hydrocarbons generated in the area.
4. Geophysical studies should be made of the region. This will provide a better structural interpretation within the region and will help select the most favorable traps and drill sites.

Most of these recommended oil and gas studies will probably be conducted, if not already completed, by industrial firms if hydrocarbon exploration continues within the area.



6.0 REFERENCES AND SELECTED BIBLIOGRAPHY

- Anderson, A.L., 1929, Geology and ore deposits of the Lava Creek district, Idaho: Idaho Bureau of Mines and Geology, Pamph. 32, 69 p.
- Anderson, A.L. and Wagne, W.R., 1946, A geological reconnaissance in the Little Wood River (Muldoon) district, Blaine County, Idaho: Idaho Bureau of Mines and Geology, Pamph. 75, 22 p.
- Armstrong, R.L., 1975, The geochronometry of Idaho: Isochron West, No. 14, 50 p.
- Armstrong, R.L., Leeman, W.P. and Malde, H.E., 1975, Dating of Quaternary and Neogene volcanic rocks of the Snake River Plain, Am. J. Sci., Vol. 275, p. 225-251.
- Armstrong, R.L., Taubeneck, W.H. and Hales, P.O., 1977, Rb/Sr and K-Ar geochronometry of the Mesozoic granitic rocks and their Sr isotopic composition, Oregon, Washington and Idaho: Geol. Soc. America Bull., Vol. 88, p. 397-411.
- Ball, L., Salisbury, J.W., Kintzinger, D.R., Veneruso, A.F. and Ward, S.H., 1979, The national geothermal exploration technology program: Geophysics, Vol. 44, p. 1721-1737.
- Bennett, E., 1980, Granitic rocks of Tertiary age in the Idaho batholith and their relation to mineralization: Econ. Geol., Vol. 75, p. 278-288.
- Beutner, E., 1977, Causes and consequences of curvature in the Sevier orogenic belt: Wyoming Geology Society Guidebook 29th Annual Field Conference, p. 353-366.
- Beutner, E. and Scholten, R., 1967, Probable Cambrian strata in east-central Idaho and their paleotectonic significance: American Assoc. of Petrol. Geologists Bull., Vol. 51, p. 2305-2311.
- Bjorlykke, A. and Sangster, D.F., 1981, An overview of sandstone lead deposits and their relationship to red-bed copper and carbonate-hosted lead-zinc deposits: Econ. Geol. 75th Anniv. Vol., p. 179-213.
- Blackstone, D., 1977, The overthrust belt salient of the Cordilleran fold belt-western Wyoming, southeastern Idaho, northeastern Utah: Joint Wyoming-Montana-Utah Geological Assoc. Guidebook, p. 367-384.
- Bollmann, D.D., 1971, Geology of east part of Mackay 3SE and west part of Mackay 4SW Quadrangles, Blaine, Butte, and Custer Counties, Idaho: Univ. of Wisconsin Milwaukee M.S. thesis, 115 p.
- Bostwick, D.A., 1955, Stratigraphy of the Wood River Formation, south-central Idaho: Journal of Paleontology, Vol. 29, No. 6, p. 941-951.



Breckenridge, R.M., 1982, Oil and gas exploration in Idaho: Idaho Bureau of Mines and Geology, Map 4, 1 sheet.

Brott, C.A., Blackwell, D.D., Mitchell, J.C., 1976, Geothermal investigations in Idaho, No. 8, Heat flow study of the Snake River Plain, Idaho: Idaho Dept. Water Resources Water Info. Bull. 30, 195 p.

Callahan, W.H., 1978, Tectonic implications of the heat flow of the western Snake River Plain, Idaho, Geol. Soc. America Bull., Vol. 89, p. 1697-1707.

Callahan, W.H., 1967, Some spatial and temporal aspects of the localization of Mississippi Valley-Appalachian-type ore deposits: Econ. Geol. Mon. 3, p. 14-19.

Carne, R., 1976, Stratabound barite- and lead-zinc-barite deposits in eastern Selwyn basin, Yukon Territory: Dept. of Indian Affairs and Northern Development, Open-File Rept. EGS 1976-14, 41 p.

Carroll, T., 1981, Memorandums: U.S. Bureau of Land Management internal files, Idaho Falls district office.

Christiansen, R.L. and McKee, E.H., 1978, Late Cenozoic volcanic and tectonic evolution of the Great Basin and Columbia-Intermontane regions: Geol. Soc. America Mem. 152, p. 283-312.

Churkin, M., Jr., 1962, Facies across Paleozoic midgeosynclinal margin of central Idaho: American Assoc. Petrol. Geologists Bull., Vol. 46, No. 5, p. 591-596.


Claypool, G.E., Love, A.H. and Maughan, E.K., 1978, Organic geochemistry, incipient metamorphism, and oil generation in black shale members of the Phosphoria Formation, western interior United States: American Assoc. Petrol. Geologists Bull., Vol. 62, p. 98-120.

Cook, E.F., 1955, Prospecting for uranium, thorium and tungsten in Idaho: Idaho Bureau of Mines and geology, Pamph. 102, 53 p.

Day, M.F., 1974, Linears map, Band 5, MSS-ERTS, northern and southern Idaho: Idaho Bureau of Mines and Geology Open-File Map, 2 Fig.

Dover, J.H., 1969, Bedrock geology of the Pioneer Mountains, Blaine and Custer Counties, central Idaho: Idaho Bureau of Mines and Geology Pamph. 142, 66 p.

Dover, J.H., 1980, Status of the Antler orogeny in central Idaho - Clarifications and constraints from the Pioneer Mountains, in Fouch, T.D. and Magethan, E.R., eds., Paleozoic Paleogeography of the West-Central United States: Soc. of Economic Paleontologists and Mineralogists, Rocky Mountain Section, West-Central United States Paleogeography Symp. 1, p. 371-386.



Dover, J.H., 1981, Chapter A-Geology of the Boulder-Pioneer study area in Mineral Resources of the Boulder-Pioneer Study Area, Blaine and Custer Counties, Idaho: U.S. Geol. Survey Bull. 1497, p. 16-75.

Dover, J.H. and Ross, R.J., Jr., 1975, Ordovician and Middle Silurian rocks in the Wildhorse window, northeastern Pioneer Mountains, central Idaho: U.S. Geological Survey Journal of Research, Vol. 3, No. 4, p. 431-436.

Ellis, A.J. and Mahon, W.A.J., 1977, Chemistry and geothermal systems: Academic Press, New York, 329 p.

Erwin, C.R., Jr., 1972, Geology of the Mackay 2SE, 2NE and part of 1NW Quadrangles, Custer County, Idaho: Wisc. Univ., Milwaukee, unpub. M.S. thesis, 128 p.

Finch, E.H., 1917, Muldoon district, Idaho: U.S. Geol. Survey Prof. Paper 97, p. 106-110.

Gutschick, R.C., Sandberg, C.A. and Sando, W.J., 1980, Mississippian shelf margin and carbonate platform from Montana to Nevada in Fouch, T.D. and Magathan, E.R. (ed.), Paleozoic Paleogeography in West-Central United States, United States Paleogeography Symp. 1: Rocky Mountain section SEPM, June 1980, p. 111-128.

Hall, W.E., Batchelder, J. and Douglass, R.C., 1974, Stratigraphic section of the Wood River Formation, Blaine County, Idaho: U.S. Geological Survey Journal of Research, Vol. 2, No. 1, p. 89-95.

Hall, W.E., Batchelder, J. and Skipp, B., 1975, Structural evolution of the Milligen and Wood River allochthons, south-central Idaho: Geol. Soc. of America Abs. with Programs, Vol. 7, No. 5, p. 611-612.

Hall, W.E., Rye, R.O. and Doe, B.R., 1978, Wood River mining district, Idaho-Intrusion related lead-silver deposits derived from country rock source: U.S. Geol. Survey, Jour. Research, Vol. 6 p. 579-592.


Haskett, G.R., 1976, Thermal springs and wells, lineaments and circular feature map of Idaho, U.S. Bureau of Rec. Map, Open-File.

Heany, M.A., 1961, Blackleaf field in Montana Oil and Gas Field Symp.: Billings Geological Society, 1961 Supplement, 3 p.

Hewitt, A.V. and Bozoin, C.N., 1962, Oxidized zinc deposits of the United States: U.S. Geol. Survey Bull. 1135.

Hobbs, S.W., 1968, Geological setting of metallic ore deposits in the northern Rocky Mountains and adjacent areas, in Ridge, J.D. (ed.), Ore Deposits in the United States 1933-1967: American Institute of Mining Metall. and Petrol. Engs., New York, p. 1351-1373.

Huh, O.K., 1967, The Mississippian System across the Wasatch line east-central Idaho, extreme southwestern Montana: Montana Geol. Society Guidebook 18th Annual Field Conf., p. 31-62.



Hunt, C.B., 1974, Natural regions of the United States and Canada: W.H. Freeman and Company, p. 409-414.

Hustedde, G.S., Copeland, J.A., Strowd, W.B., Mitchell, V.E. and Bennett E.H., 1981, Mines and prospects of the Hailey Quadrangle, Idaho: Idaho Bur. Mines and Geology, Mines and Prospects Map Series, 41 p.

King, P.B., 1969, The tectonics of North America - a discussion to accompany the tectonic map of North America, scale 1:5,000,000: U.S. Geol. Survey Prof. Paper 628, 95 p.

Kuntz, M.A., Skipp, B., Embree, G.F., Jr., Hogan, E.D. and Williams, E.J., 1979, Geologic map of the Lava Ridge-Hell's Half Acre, eastern Snake River Plain, Idaho: U.S. Geol. Survey Open-File Rept. 669.

Lamb, C.F., 1980, Painter reservoir field -- giant in Wyoming Thrust belt: American Assoc. Petrol. Geologists Bull., Vol. 64, No. 5, p. 638-644.

Loucks, G.H., 1977, Geologic history of the Devonian, northern Alberta to southwest Arizona: Wyoming Geol. Assoc. Guidebook, 29th Annual Field Conf., p. 119-134.

Lukowicz, L.J., 1971, Geology of Mackay 3SW and west part of Mackay 3SE Quadrangle, Blaine County, Idaho: Univ. of Wisconsin-Milwaukee M.S. thesis, 127 p.

Mabey, D.R., 1981, Chapter B-Aeromagnetic studies of the Boulder-Pioneer wilderness study area, Blaine and Custer Counties, Idaho in Boulder-Pioneer Study Area, Idaho: U.S. Geol. Survey Bull. 1497, p. 77-84.

Maley, T., 1983, Handbook of mineral law: MMRC Publ., Boise, Idaho 239 p.


Mamet, B.L., Skipp, B.A.L., Sando, W.J. and Mapel, W.J., 1971, Biostratigraphy of Upper Mississippian and associated Carboniferous rocks in south-central Idaho: American Assoc. Petrol. Geologists Bull., Vol. 55, No. 1, p. 20-33.

Mansfield, G.R., 1927, Geography, geology and mineral resources of part of southeastern Idaho: U.S. Geol. Survey Prof. Paper 152, 453 p.


Mapel, W.J. and Sandberg, C.A., 1968, Devonian paleotectonics in east-central Idaho and southwestern Montana: U.S. Geol. Survey Prof. Paper 600D, p. D116-D125.

McCaslin, J.C., 1980, Amoco scores deep Nugget hit in overthrust: Oil and Gas Journal, Vol. 78, No. 33, p. 161.

_____, 1981, Six new fields make list of U.S. oil giants: Oil and Gas Journal, Vol. 79, No. 4, p. 153.



- McIntyre, D.H., Hobbs, S.W., Marvin, R.F. and Mehnert, H.H., 1976, Late Cretaceous and Eocene ages for hydrothermal alteration and mineralization, Bayhorse district and vicinity, Custer County, Idaho: *Isochron*/west, No. 16, p. 11-12.
- Mitchell, A.H.G. and Garson, M.S., 1981, Mineral deposits and global tectonic settings: Academic Press, London, 405 p.
- Mitchell, J.C., Johnson, L.L. and Anderson, J.E., 1980, Geothermal investigations in Idaho: Part 9, Potential for direct heat application of geothermal resources: Idaho Dept. of Water Resources, Water Inf. Bull. 30, 396 p.
- Mohley, L.E., 1971, Petroleum potential of Idaho-Wyoming overthrust belt in Future Petroleum Provinces of the United States -- Their Geology and Potential, I.H. Cram (ed.): American Assoc. Petrol. Geologists, Mem. 9, Vol. 1, p. 509-529.
- Muffler, L.J.P. (ed.), 1979, Assessment of geothermal resources of the United States, 1978: U.S. Geol. Survey Circ. 790, 163 p.
- Nelson, W.H. and Röss, C.P., 1969, Geologic map of the Mackay Quadrangle, south-central Idaho: U.S. Geol. Survey, Misc. Geol. Map I-580.
- Nilsen, T.H., 1977, Paleogeography of Mississippian turbidites in south-central Idaho, in Stewart, J.H., Stevens, C.H. and Frische, A.L., Paleozoic Paleogeography of the Western United States: Society of Econ. Paleontologists and Mineralogists, Pacific Coast Paleogeography Sym. 1, p. 275-297.
- Noble, J.A., 1970, Metal provinces in the western United States: Geol. Soc. America Bull., Vol. 81, p. 1607-1624.
- _____, 1974, Metal provinces and metal finding in the western United States: Mineral. Deposit, Vol. 9, p. 1-25.
- Pankratz, L.W. and Ackermann, H.D., 1982, Structure along the northwest edge of the Snake River Plain interpreted from seismic refraction: Journ. Geophys. Res., Vol. 87, No. B4, p. 2676-2682.
- Pauli, R.A., 1974, Upper Cretaceous? to lower Eocene post-orogenic conglomerate, south-central Idaho: Geol. Soc. of America Abs. with Programs, Vol. 6, No. 5, p. 466.
- Pauli, R.A. and Gruber, D.P., 1977, Little Copper Formation - New name for lowest formation of Mississippian Copper Basin Group, Pioneer Mountains, south-central Idaho: American Assoc. Petrol. Geologists Bull., Vol. 61, No. 2, p. 256-262.
- Pauli, R.A., Wolbrink, M.A., Wolkman, R.G. and Grover, R.L., 1972, Stratigraphy of Copper Basin Group, Pioneer Mountains, south-central Idaho: American Assoc. Petrol. Geologists Bull., Vol. 56, No. 8, p. 1370-1401.



Perry, W.J., Ryder, R.T. and Maughan, E.K., 1981, The southern part of the southwest Montana Thrust belt: a preliminary re-evaluation of structure, thermal maturation and petroleum potential: Montana Geol. Society 1981 Field Conf. and Symp. Guidebook to Southwest p. 261-273.

Peterson, J.A., 1981, General stratigraphy and regional paleostructure of the western Montana Thrust belt: Montana Geol. Society Field Conf. and Symp. Guidebook to Southwest Montana, p. 5-35.

Poole, F.G., 1974b, Antler flysch in central Idaho: U.S. Geol. Survey Prof. Paper 900, p. 34.

Poole, F.G. and Sandberg, C.A., 1977, Mississippian paleogeography and tectonics of the western United States in Stewart, J.H., Stevens, C.H. and Fritsche, A.E., eds., Paleozoic Paleogeography of the Western United States: Soc. Econ. Paleontologists and Mineralogists, Pacific Section, Pacific Coast Paleogeography Symp. 1, p. 67-85.

Rember, W.C. and Bennett, E.H., 1979a, Geologic map of the Hailey Quadrangle, Idaho: Idaho Bureau of Mines and Geology Geologic Map Series Hailey 2° Quadrangle, 1 sheet.

_____, 1979b, Geologic map of the Idaho Falls Quadrangle, Idaho: Idaho Bureau of Mines and Geology Geologic Map Series Twin Falls 2° Quadrangle, 1 sheet.

Rooney, L.F., 1956, Organic carbon in Phosphoria Formation: American Assoc. Petrol. Geologists Bull., Vol. 40, p. 2267-2271.

Ross, C.P., 1931, A classification of lode deposits of south-central Idaho: Econ. Geol., Vol. 26, p. 169-185.

_____, 1937, Geology and ore deposits of the Bayhorse region, Custer County, Idaho: U.S. Geol. Survey Bull. 877, 161 p.

_____, 1962a, Paleozoic seas of central Idaho: Geol. Soc. of America Bull., Vol. 73, No. 6, p. 769-794.


_____, 1962b, Stratified rocks in south-central Idaho: Idaho Bureau of Mines and Geology Pamph. 125, 126 p.

Ross, S.H., 1971, Geothermal potential of Idaho: Idaho Bureau of Mines and Geology, Pamph. 150, 72 p.

Rosta, J., 1982, Silver in 1981-1982, 113th Annual Survey of Mineral Commodities: Engineering and Mining Journal, March 1982, p. 143-145.

Ruppel, E.T., 1964, Strike-slip faulting and broken basin-range in east-central Idaho and adjacent Montana: in Geological Survey Research 1964: U.S. Geol. Survey Prof. Paper 501-C, p. C14-C18.

_____, 1978, The Medicine Lodge thrust system, east-central Idaho and southwest Montana: U.S. Geol. Survey Prof. Paper 1031, 23 p.



Ruppel, E.T., 1982, Cenozoic block uplifts in east-central Idaho and southwest Montana: U.S. Geol. Survey Prof. Paper 1224, 24 p.

Ruppel, E.T., Wallace, C.A., Schmidt, R.G. and Lopez, D.A., 1981, Preliminary interpretation of the Thrust belt in southwest and west-central Montana and east-central Idaho: Montana Geol. Society Field Conf., p. 139-159.

Ruppel, E.T., Watts, K.C. and Peterson, D.L., 1970, Geologic, geochemical and geophysical investigations in the northern part of the Gilmore mining district, Lemhi County, Idaho: U.S. Geol. Survey Open-File Rept., 56 p.

Sandberg, C.A., 1975, McGowan Creek Formation, new name for Lower Mississippian flysch sequence in east-central Idaho: U.S. Geol. Survey Bull. 1405-E, 11 p.

Sandberg, C.A., Hall, W.E., Batchelder, J.N. and Axelsen, C., 1975, Stratigraphy, conodont dating and paleotectonic interpretation of the type Milligen Formation (Devonian), Wood River area, Idaho: U.S. Geol. Survey Journal Research, Vol. 3, No. 5, p. 707-720.

Sandberg, C.A., Mapel, W.J. and Huddle, J.W., 1967, Age and regional significance of basal part of Milligen Formation, Lost River Range, Idaho: U.S. Geol. Survey Research 1967: U.S. Geol. Survey Prof. Paper 11576-C, p. C127-C131.

Savage, C.N., 1969, Distribution and occurrence of identified oil and gas rocks, Idaho Bureau of Mines and Geology Bull. 23, 99 p.

Scholten, R., 1957, Paleozoic evolution of the geosynclinal margin north of the Snake River Plain, Idaho-Montana: Geol. Soc. of America Bull., Vol. 68, p. 151-170.


_____, 1967, Structural framework and oil potential of extreme southwestern Montana: Montana Geol. Society Guidebook 18th Annual Field Conf., p. 7-20.

_____, 1968, Model for evolution of Rocky Mountains east of Idaho batholith: Tectonophysics, Vol. 16, No. 2, p. 109-126.

_____, 1973, Gravitational mechanism in the northern Rocky Mountains of the United States in DeLong, K.A. and Scholten, R. (eds.), Gravity and Tectonics: New York, John Wiley and Sons, 1973, p. 473-489.

Scholten, R. and Hait, M.H., Jr., 1962, Devonian System for shelf edge to geosyncline, southwestern Montana-central Idaho in Three Forks Belt Mountain Area, and Symp., The Devonian system of Montana and Adjacent Areas, Billings Geol. Soc., p. 13-22.

Simons, F.S., 1981, Chapter C-A geological and geochemical evolution of the mineral resources of the Boulder-Pioneer wilderness study area, Blaine and Custer Counties, Idaho in Boulder-Pioneer Study Area, Idaho: U.S. Geol. Survey Bull. 1497, p. 86-180.



Singer, D.A. and Mosier, D.L., 1981, A review of regional mineral resource assessment methods: *Econ. Geol.*, Vol. 76, No. 5, p. 1006-1015.

Skipp, B.A.L., 1961, Stratigraphic distribution of endothyrid Foraminifera in Carboniferous rocks of the Mackay Quadrangle, Idaho in *Geological Survey Research 1961: U.S. Geol. Survey Prof. Paper 424-C*, p. C239-C244.

_____, 1981, A synopsis of the structure and stratigraphy of south-central Idaho -- Contributions by the U.S. Geological Survey since 1975: *Montana Geol. Society Field Conf. Southwest Montana*, p. 365-372.

Skipp B. and Hait, M.H., Jr., 1977, Allochthons along the northeast margin of the Snake River Plain, Idaho: *Wyoming Geol. Assoc. Guidebook 29th Annual Field Conf.*, p. 499-515.

Skipp, B. and Hall, W.E., 1980, Upper Paleozoic paleotectonics and paleogeography of Idaho in Fouch, T.D. and Magathan, E.R., eds., *Paleozoic Paleogeography of the West-Central United States: Soc. of Economic Paleontologists and Mineralogists, Rocky Mountain Section, West-Central United States Paleogeography Symp. 1*, p. 387-422.

_____, 1975, Structure and Paleozoic stratigraphy of a complex of thrust plates in the Fish Creek Reservoir area, south-central Idaho: *U.S. Geol. Survey Journal of Research*, Vol. 3, No. 6, p. 671-689.

Skipp, B. and Sandberg, B.A., 1975, Silurian and Devonian miogeosynclinal and transitional rocks of the Fish Creek Reservoir window, central Idaho: *U.S. Geol. Survey Journal of Research*, Vol. 3, No. 6, p. 691-706.


Skipp, B., Sando, W.J., and Hall, W.E., 1979, The Mississippian and Pennsylvanian (Carboniferous) Systems in the United States -- Idaho: *U.S. Geol. Survey Prof. Paper 1010, part AA*, p. 1-42.

Skipp, B., Hoggan, E.D., Schleicher, D.L. and Douglass, R.C., 1979, Upper Paleozoic carbonate bank in east-central Idaho-Snaky Canyon, Bluebird Mountain and Arco Hills Formations and their paleotectonic significance: *U.S. Geol. Survey Bull.* 1486, 78 p.


Sloss, L.L., 1954, The Lemhi arch, a mid-Paleozoic positive element in south-central Idaho: *Geol. Soc. America Bull.*, Vol. 65, p. 365-368.

Smith, R.B., 1978, Seismicity, crustal structure and intraplate tectonics of the interior of the western Cordillera: *Geol. Soc. America Mem.* 512, p. 111-144.

Smith, R.B. and Sbar, M., 1974, Contemporary tectonics and seismicity of the western United States with emphasis on the Intermountain seismic belt: *Geol. Soc. America Bull.*, Vol. 85, p. 1205-1218.




- Strowd, W.B. Mitchell, V.E., Hustedde, G.S. and Bennett, E.H., 1981, Mines and Prospects of the Idaho Falls Quadrangle, Idaho: Idaho Bureau of Mines and Geology, Mines and Prospects Map Series, 10 p.
- Swanberg, C.A. and Blackwell, D.D., 1973, Areal distribution and geophysical significance of heat generation in the Idaho batholith and adjacent intrusions in eastern Oregon and western Montana: Geol. Soc. America Bull., Vol. 84, p. 1261-1282.
- Thomasson, M.R., 1959, Late Paleozoic stratigraphy and paleotectonics of central and eastern Idaho: Wisconsin Univ., Madison, Ph.D. thesis, 244 p.
- Tuchek, E.T. and Ridenour, J., 1981, Chapter D-Economic appraisal of the Boulder-Pioneer wilderness study area, Blaine and Custer Counties, Idaho in Boulder-Pioneer Study Area, Idaho: U.S. Geol. Survey Bull. 1497, p. 181-292.
- Umpleby, J.B., 1917, Geology and ore deposits of the Mackay region, Idaho: U.S. Geol. Survey Prof. Paper 97, 129 p.
- Umpleby, J.B., Westgate L.G. and Ross, C.P., 1930, Geology and ore deposits of the Wood River region, Idaho, with a description of The Minnie Moore and nearby mines by D.F. Hewett: U.S. Geol. Survey Bull. 814, 250 p.
- Volkman, R.G., 1971, Geology of the Mackay 3NE Quadrangle, Blaine and Butte Counties, Idaho: Univ. of Wisconsin-Milwaukee M.S. thesis, 121 p.
- Wodzicki, A. and Krason, J., 1981, National uranium resource evaluation, Dubois Quadrangle, Idaho and Montana: U.S. Dept. of Energy, Rept. GJQ-008(81), 67 p.
- Wolbrink, M.A., 1970, Geology of the Mackay 3NW Quadrangle, Custer and Blaine Counties, Idaho: Univ. of Wisconsin-Milwaukee M.S. thesis, 108 p.



APPENDIX I

WILDERNESS STUDY AREA MAPS



53-1
LITTLE WOOD RIVER

53-5
FRIEDMAN CREEK

R. 20 E.

R. 21 E.

R. 22 E.


52-14
LITTLE WOOD RIVER

53-5
FRIEDMAN CREEK



R. 20 E.


R. 21 E.

R. 22 E.




LEGEND






31-14
APPENDICITIS HILL






LEGEND

-  WILDERNESS STUDY AREA
 AREA WITHOUT CHARACTERISTICS




31-17
WHITE KNOB MOUNTAINS





APPENDIX II

FIELD VISIT REPORT



To: Pioneer Mountains GRA File
Fm: Greg Fernetto, Bill Jones

Subject: Visit to GRA - October 12 and 13, 1982


The Muldoon mine in Muldoon Canyon was visited since similar mineralization may exist in the nearby Friedman Creek WSA. The WSA was also visited and work performed along with significant observations are summarized below:

Muldoon Mine

Mineralization at the mine consists of coarse pyrite, sphalerite and some chalcopyrite and specular hematite. The sulfides occur in veins and breccia zones associated with altered hypabyssal(?) porphyritic dacite dikes which have intruded along a fault zone trending about N50W. The country rocks are quartzites with lesser conglomerate and shale. The trend of the mineralized zone projects into the Friedman Creek WSA.


Friedman Creek WSA

Traverses were made up of the drainages and along some ridges in the WSA to examine the geology and collect stream sediment samples. Thirty-three samples were collected. These samples were sent to Rainbow Resource Laboratories in Anchorage for analysis. The WSA is underlain mainly by quartzites with some shale and conglomerate, capped by Challis Volcanics. Altered dikes and mineralized talus similar to the rocks at the Muldoon mine were found in several places.



APPENDIX III

ANALYTICAL DATA FOR GEOCHEMICAL SAMPLES COLLECTED IN THE
FRIEDMAN CREEK WSA



Sample No.	Sample Type	Cu (ppm)	Pb (ppm)	Zn (ppm)	Ag (ppm)	Au (ppm)
28013	SS	40	95	135	1.0	.04
28014	SS	55	45	345	4.0	.02
28015	SS	65	50	195	2.2	.02
28016	SS	95	50	135	2.2	.02
28017	SS	40	65	110	.4	.02
28018	SS	35	30	80	.2	.04
28019	SS	45	45	150	.2	.04
28020	SS	25	40	75	.2	.10
28030	SS	40	35	165	1.2	.04
28031	SS	45	45	390	.2	.04
28032	SS	90	60	500	1.2	.04
28033	SS	35	40	230	1.0	.02
28034	SS	30	40	135	.6	.02
28035	SS	30	45	255	.6	.02
28036	SS	335	335	450	5.0	.10
28037	SS	45	55	195	1.0	.04
28041	SS	40	175	145	1.0	.02
28042	SS	30	50	230	.6	.02
28043	SS	45	45	100	1.2	.02
28044	SS	60	55	465	1.0	.04
28045	SS	30	50	280	.2	.02
28046	SS	40	55	175	.2	.02
28047	SS	35	50	140	.4	.04
28048	SS	35	50	175	.2	.02
28049	SS	30	30	90	1.2	.02
28050	SS	40	45	110	1.2	.02
28051	SS	95	50	100	.6	.02
28052	SS	50	170	140	.8	.02
28053	SS	30	75	100	.2	.02
28054	SS	40	40	90	.2	.02
28055	SS	45	35	465	.4	.02
28056	SS	140	60	150	1.0	.02
28057	SS	45	40	400	.4	.02

Atomic absorption analyses by Rainbow Resource Laboratories, Anchorage, Alaska.

ELM Library
D-553A, Building 50
Denver Federal Center
P. O. Box 25047
Denver, CO 80225-0047

1944-1945
1946-1947
1948-1949
1950-1951
1952-1953
1954-1955
1956-1957
1958-1959
1960-1961
1962-1963
1964-1965
1966-1967
1968-1969
1970-1971
1972-1973
1974-1975
1976-1977
1978-1979
1980-1981
1982-1983
1984-1985
1986-1987
1988-1989
1990-1991
1992-1993
1994-1995
1996-1997
1998-1999
2000-2001
2002-2003
2004-2005
2006-2007
2008-2009
2010-2011
2012-2013
2014-2015
2016-2017
2018-2019
2020-2021
2022-2023
2024-2025

1944-1945
1946-1947
1948-1949
1950-1951
1952-1953
1954-1955
1956-1957
1958-1959
1960-1961
1962-1963
1964-1965
1966-1967
1968-1969
1970-1971
1972-1973
1974-1975
1976-1977
1978-1979
1980-1981
1982-1983
1984-1985
1986-1987
1988-1989
1990-1991
1992-1993
1994-1995
1996-1997
1998-1999
2000-2001
2002-2003
2004-2005
2006-2007
2008-2009
2010-2011
2012-2013
2014-2015
2016-2017
2018-2019
2020-2021
2022-2023
2024-2025